




UKRAINE
 NATIONAL UNIVERSITY OF LIFE AND ENVIRONMENTAL
 SCIENCES OF UKRAINE
 Department of Analytical and Bioinorganic Chemistry & Water
 Quality

WORKBOOK

in CHEMISTRY (V) (analytical)

of student _____
 (Name, Surname)

(group, year of study)
 Branch of knowledge – 10 Natural Sciences
 Speciality – 101 Ecology



pH Indicator Chart

Kyiv-2019

УДК 546 (07)

Робочий зошит призначено для виконання лабораторного практикуму з аналітичної хімії. Включає методики виконання експериментальних задач, форми протоколів аналізу, питання для самоперевірки.

Рекомендовано методичною радою факультету захисту рослин, біотехнологій та екології Національного університету біоресурсів і природокористування України для спеціальності 101 – Екологія, для яких викладання ряду дисциплін проводиться англійською мовою (протокол № ____ від _____ 2019 р.).

Рецензенти: д.х.н., проф. Максим В.І., НУБіП України;
к.х.н., доц. Бухтіяров В.К., НУБіП України.

Voitenko, L.V. Workbook in analytical (V) chemistry (Speciality 101-Ecology) / Kyiv : DDP “Expo-druk”. – 2019. – 92 p.

Відповідальний за випуск: к.х.н., доц. Войтенко Л.В.

UDC 546 (07)

The workbook is intended to train in lab course of Analytical chemistry. It contains the description of experimental strategies, lab techniques, the templates of lab reports, test questions for own training.

Recommended for publication by the by the Methodical board of the Plant Protection, the Biotechnologies & Ecology faculty of National University of Life and Environmental Sciences, for Speciality 101 – Ecology.

Report # ____, Data _____ 2019.

Reviewers: Doctor of Chem. Sci., Prof. Maksin V.I., NUBiP of Ukraine;
Candidate of Chem. Sci., Ass. Prof. Buchtiyarov V.K., NUBiP of Ukraine.

Responsible for the edition: Cand. of Chem. Sci., Ass. Prof. Voitenko L.V.

CONTENT

INTRODUCTION.....	5
GENERAL CHEMISTRY SAFETY AND LABORATORY RULES.....	5
1.2. LAB GLASSWEARE CLEANING RULES	6
CHAPTER 1. QUALITATIVE ANALYSIS	8
1.1. ANALYTICAL REAGENTS AND REACTIONS.....	8
1.2. QUALITATIVE REACTIONS OF THE CATIONS.....	9
1.3. THE FIRST GROUP OF CATIONS.....	11
1.3.1. Characteristic reactions of Ammonia cations NH_4^+	11
1.3.2. Characteristic reactions of Potassium cations K^+	12
1.3.3. Characteristic reactions of Sodium cations Na^+	13
1.4. THE SECOND GROUP OF CATIONS.....	14
1.4.1. Action of group reagent $(\text{NH}_4)_2\text{HPO}_4$	14
1.4.2. Characteristic reactions of Magnesium cations Mg^{2+}	16
1.4.3. Characteristic reactions of Calcium cations Ca^{2+}	16
1.4.4. Characteristic reactions of Strontium cations Sr^{2+}	17
1.4.5. Characteristic reactions of Manganese cations Mn^{2+}	18
1.4.6. Characteristic reactions of Ferrous cations Fe^{2+}	19
1.4.7. Characteristic reactions of Ferric cations Fe^{3+}	20
1.4.8. Characteristic reactions of Aluminium cations Al^{3+}	21
1.5. THE THIRD GROUP OF CATIONS	22
1.5.1. Action of group reagent $(\text{NH}_4)_2\text{HPO}_4$	22
1.5.2. Characteristic reactions of Copper cations Cu^{2+}	23
1.5.4. Characteristic reactions of Cobalt cations Co^{2+}	26
1.5.5. Characteristic reactions of Nickel cations Ni^{2+}	28
1.6. THE FOURTH GROUP OF CATIONS	29
1.6.1. Action of group reagent HCl	29
1.6.2. Characteristic reactions of Silver cations Ag^+	30
1.6.3. Characteristic reactions of Lead cations Pb^{2+}	31
1.7. REACTIONS OF THE ANIONS	33
1.7.1. The first group of anions.....	33
1.7.2. The second group of anions.....	42
1.7.3. The third group of anions.....	44
1.8. PROTOCOL CATIONS MIXTURE OF ANALYSIS.....	47
1.9. QUALITATIVE ANALYSIS OF THE INORGANIC SUBSTANCES	59
1.9.1 A soluble salt.....	59
1.9.2 Insoluble substances.....	66
EXAMPLES OF CONTROL TASKS FOR QUALITATIVE ANALYSIS	69

CHAPTER 2. QUNTITATIVE ANALYSIS	71
2.1. Lab work - Precipitate gravimetry (mass analysis). Determination of Barium content in Hydrated Barium Chloride.....	71
2.2. Laboratory work – Volumetric neutralization method. Determination of alkali concentration in solution.....	73
2.3. Lab work – Oxidation-reduction (RedOx) titration (RedOxmetry). Determination of Iron(II) content in Mohr’s salt.....	77
2.4. Laboratory work – Iodometry. Determination of Copper(II) Content in Copper Vitriol	81
2.5. Laboratory work – Complexonometry. Determination of Calcium concentration in solution	83
EXAMPLES OF CONTROL TESTS FOR QUANTITATIVE ANALYSIS	87
MY RATING TABLE	92

INTRODUCTION

GENERAL CHEMISTRY SAFETY AND LABORATORY RULES

Chemistry laboratories can be hazardous if the rules are not followed. During a chemistry course a student may handle materials which are carcinogenic, poisonous, flammable, and explosive. Some of these materials and equipment may also cause severe burns, cuts, or bruises if handled improperly or carelessly. Most accidents that occur in the chemistry laboratory are a result of carelessness, impatience, improper or unauthorized experimentation, and disregard for safety rules or proper operating procedures. In order to minimize the chances of an accident in the laboratory certain rules and regulations must be obeyed at all times when one is working or observing in a chemical laboratory. Therefore, it is not advisable for anyone to work in a laboratory without proper knowledge of the dangers involved. Due to the inherent dangers present in a chemical laboratory exercise, it should be understood that the following rules must be obeyed to minimize the chance of an accident. The student is expected to exercise proper judgement and extreme caution at all times when working in the laboratory.

1. Do not perform unauthorized experiments or work in a laboratory alone.

2. Approved eye protection must be worn at all times in the laboratory. If you do get a chemical in your eye rinse immediately with large quantities of water using the eye-wash stations.

3. Long hair and loose clothing must be confined while in a laboratory.

4. Appropriate clothing must be worn at all times while in the laboratory. Your legs must be completely covered below the knee by your choice of clothing. If your clothing does not meet the requirement you may choose to wear an approved laboratory coat or apron which does cover your legs to your knees.

5. Closed shoes with socks must be worn.

6. Know the location and proper use of fire extinguishers, fire blankets, eye wash devices, and first aid kits.

7. Before obtaining any chemicals carefully read the label on the reagent bottles.

8. Eating, smoking, and drinking are not allowed in a chem laboratory.

9. Thoroughly wash your hands after leaving the laboratory.

10. Use the fume hoods when toxic or irritating vapours are involved.

11. Mouth suction is never used to fill a pipette.

13. Never direct the open end of test tube toward yourself or anyone else.

14. Never pour water into concentrated acid (**especially H₂SO₄**).
15. Learn the proper procedure for igniting and operating a laboratory burner. Always extinguish the flame when the burner is not being used. Make sure that all flammable reagents are well removed before lighting the burner.
16. Liquid and solid waste containers must be properly used at all times.
17. Never place chemicals directly on the balance pan. Always use a proper weighing container when using a balance to weigh a chemical. Never pour chemicals directly over the balance.
18. Never return unused chemicals to their original container (unless directed to do so by the instructor).
19. Securely replace lids, caps, and stoppers after removing reagents from containers.
20. Always wipe spatulas clean before and after inserting into reagent bottles.
21. Report any accident and/or injury, however minor, to your instructor immediately.
22. Never place anything that is not directly required for the experiment on laboratory desks; other items may interfere with the experiment.
23. All personal belongings should be placed in the bookcases as you enter the laboratory.
24. Clean up any spill immediately.
25. Before leaving the laboratory, make sure your work area is clean and dry.
26. Your instructor is available for any assistance you may need. Never hesitate to ask questions especially if there is any question concerning proper operating procedure. Be sure that you understand every instruction before proceeding.

1.2. LAB GLASSWEARE CLEANING RULES

In the analysis of samples the preparation of scrupulously clean glassware is mandatory. Lab glassware cleaning procedures must follow specific method requirements. If procedures are not listed then the method of cleaning should be adapted to both the substances that are to be removed, and the determinations (tests) to be performed. Recommendations for such cleaning procedures are listed below.

Analysis/Parameter

Cleaning Procedure
(In order specified)

ORGANICS

INORGANICS

Trace Metals: 1-4, 9, 8 (optional), 4

Nutrients, Minerals: 1-4, 8, 4

Solids: 1-4, 11
(Volatile Solids 16)

Non-Metals, Physical Properties:
(Cyanide) 1-4, 14

MICROBIOLOGY 1-4 (Sterilize per approved method)

BIOASSAY

Freshwater: 18, 2, 3, 9 or 8, 4, 5, 4, 20

Marine & Estuarine: 19, 2, 3, 9 or 8, 4, 5, 4, 20

RADIONUCLIDES 17, 3, 8, 4

Cleaning Procedures:

1. Remove all labels using sponge or acetone.
2. Wash with hot tap water and a brush to scrub inside of glassware, stopcocks, and other small pieces, if possible, using a suitable laboratory-grade detergent. Organics - Liquinox, Alconox or equivalents. Inorganic anions - Liquinox or equivalent. Inorganic cations - Liquinox, Acationox, Micro or equivalents.
3. Rinse thoroughly with hot tap water.
4. Rinse thoroughly with distilled water (DW).
5. Rinse thoroughly with pesticide grade Acetone.
6. Rinse thoroughly with pesticide grade Methanol.
7. Rinse thoroughly with pesticide grade Hexane.
8. Rinse or soak with 1:1 HCl (Hydrochloric Acid).
9. Rinse or soak with >10% HNO₃ (Nitrate Acid).
10. Bake at 105 °C for 1 hour.
11. Bake at 180 °C (prior to use as per method).
12. Drain, then heat in muffle furnace for 30-60 minutes at 400 °C.
13. Clean, dry glassware should be sealed and stored in dust-free environment.
14. Soak in oxidizing agent (Chromic acid or equivalent); preferably hot (40-50 °C).
15. Last step (prior to use) should be a rinse with the solvent used in analysis.
16. Drain, then heat in muffle furnace for 1 hour at 550 °C.
17. Heat 1 hour in EDTA solution at 90-100 °C.
18. New glassware must be soaked overnight in 10% HNO₃ or HCl.
19. New glassware must be soaked overnight in seawater.
20. Rinse thoroughly with DW.

CHAPTER 1. QUALITATIVE ANALYSIS

1.1. ANALYTICAL REAGENTS AND REACTIONS

Using chemical method the analyzed substance is transformed into new compound having characteristic properties: colour, odour, and defined physical state. Chemical transformation which takes place is called qualitative analytical reaction and the substances initiated this transformation are called **analytical reagents**.

Group reagents react with the group of ions and provide the same effect; **specific reagents** react with only one ion; **selective reagents** react with a few ions.

Analytical reactions to be carried out are mostly influenced by the following:

- pH;
- Ion concentration;
- Temperature.

The sensitivity of the reaction is determined by the following:

- Detectable minimum;
- Minimum concentration;
- Maximum dilution.

Detectable minimum (m) is the less quantity of substance or ion, which can be detected by given reaction at certain conditions. It is expressed in $\gamma = 10^{-6}$ g.

Concentration limit (c) is the minimum concentration of determined ion in solution at which the analytical reaction gives positive effect. Concentration limit is expressed as the ratio of substance mass (1 g) to solution volume (in mL). For example, for qualitative determination of Cu^{2+} by reaction with water solution of ammonia $\text{NH}_3 \cdot \text{H}_2\text{O}$ minimum concentration is 1 : 250 000. It means that 1 g of Cu^{2+} may be detected in 250 liters of analyzing solution. Analytical tests must have **not less than 1 : 1000**.

Maximum dilution (MD) is an index inverse to minimum concentration: $\text{MD} = 1/c$. Maximum dilution indicates at which solution volume 1 g of the substance should be presented in order the reaction to be visible.

So the sensitivity of reaction is higher if maximum dilution is bigger; but detectable minimum and minimum concentration are is smaller.

Detectable minimum and concentration limit are interrelated:

$$m = c \cdot V \cdot 10^6;$$

$$c = m/V \cdot 10^6, \text{ where}$$

V - volume of solution, mL.

1.2. QUALITATIVE REACTIONS OF THE CATIONS


Ammonia-phosphate classification of cations is based (table 1) on the difference in solubility of phosphates in water, weak and strong acids, alkali and ammonia.

Environmental practice is connected with studying of the next common biologically active cations:

- Macronutrients: NH_4^+ , Ca^{2+} , K^+ , Na^+ , Mg^{2+} ;
- Micronutrients: Mn^{2+} , Zn^{2+} , Co^{2+} , Ni^{2+} , Cu^{2+} , Ag^+ , Fe^{2+} , Fe^{3+} ;
- Cations of toxic and radioactive elements: Al^{3+} ; Sr^{2+} , Cu^{2+} , Ag^+ , Pb^{2+} .

The main idea is to use the separation of cations analyzing the mixture.


For example, **propose** the order of separation of cations in mixture.




Answer:

1. Cation Co^{2+} is included to the 3d group, Ca^{2+} – 2^d group, A-subgroup (see Table 1).

2. The 2^d and 3^d cations group reagent is the same in fact – $(\text{NH}_4)_2\text{HPO}_4$ (see Table 1). So, adding this reagent, both cations of mixture were precipitated:



3. For separating of sediments obtained we use their different solubility in excess of water ammonia $\text{NH}_3 \cdot \text{H}_2\text{O}$: sediments of the 2^d cations group are insoluble, the 3^d one – soluble:



So, the chart of operation is:




Table 1 – Principles of ammino-phosphate classification of cations

Group	Cations	Group reagent	Properties of sediments				
			Solubility in:				
			H ₂ O	NH ₃ ·H ₂ O	CH ₃ CO -OH	HCl, HNO ₃	NaOH
1	NH ₄ ⁺ , Na ⁺ , K ⁺	no	-	-	-	-	-
2	A Mg ²⁺ , Mn ²⁺ , Ca ²⁺ , Sr ²⁺ , Fe ²⁺	(NH ₄) ₂ HPO ₄ + NH ₃ ·H ₂ O	Insoluble in water	Insoluble	Soluble	Soluble	Insoluble
	B Fe ³⁺ , Al ³⁺				Insoluble		
3	Cu ²⁺ , Zn ²⁺ , Co ²⁺ , Ni ²⁺	(NH ₄) ₂ HPO ₄ or Na ₂ HPO ₄	Insoluble in water	[Me ⁿ⁺ (NH ₃) _{(2-3)n}] ⁿ⁺ soluble	Soluble	Soluble	Only Zn ₃ (PO ₄) ₂ soluble, forming Na ₂ [Zn(OH) ₄]
4	Ag ⁺ , Pb ²⁺	HCl	Insoluble (PbCl ₂ soluble partially in boiling H ₂ O)	[Ag(NH ₃) ₂]Cl soluble	Insoluble	Insoluble	AgCl – insoluble; PbCl ₂ soluble, forming Na ₂ [Pb(OH) ₄]

1.3. THE FIRST GROUP OF CATIONS

1.3.1. Characteristic reactions of Ammonia cations NH_4^+

Content	Description	Observation	Explanation
<p>1. Alkali (NaOH, KOH) at heating. Reaction is <i>highly sensitive, specific; concentration limit</i> $c=1:5000000$.</p>	Add 4-5 drops of NaOH or KOH solution to 3-4 drops of ammonia salt and heat in a water bath. Gas may be identified by its odour (cautiously smell the vapor after moving the test-tube from the water bath), or by its turning moistened litmus paper blue.		$\text{NH}_4^+ + \text{OH}^- = \text{NH}_3 + \text{H}_2\text{O}$ $\text{NH}_4\text{Cl} + \text{NaOH} \longrightarrow \text{NH}_3\uparrow + \text{NaCl} + \text{H}_2\text{O}.$
<p>2. Nessler's reagent $\text{K}_2[\text{HgI}_4] + \text{KOH}$ (alkaline solution of potassium tetraiodomercurate(II)). Reaction is <i>extremely sensitive, concentration limit</i> $c = 1:10000000$. Interferences cations formed insoluble hydroxides (Fe^{2+}, Fe^{3+}, Cu^{2+} et al.)</p>	Add 2-3 drops of Nessler's reagent to 1 drop of any soluble ammonia salt.		$\text{NH}_4^+ + 2[\text{HgI}_4]^{2-} + 4\text{KOH} =$ $\left[\begin{array}{c} \text{Hg} \\ \diagdown \quad \diagup \\ \text{O} \quad \text{NH}_2 \\ \diagup \quad \diagdown \\ \text{Hg} \end{array} \right] \text{I} \downarrow + 7 \text{I}^- + 3\text{H}_2\text{O};$ $\text{NH}_4\text{Cl} + 2\text{K}_2[\text{HgI}_4] + 4\text{KOH} =$ $\left[\begin{array}{c} \text{Hg} \\ \diagdown \quad \diagup \\ \text{O} \quad \text{NH}_2 \\ \diagup \quad \diagdown \\ \text{Hg} \end{array} \right] \text{I} \downarrow + 7\text{KI} +$ $\text{KCl} + 3\text{H}_2\text{O}.$

1.3.2. Characteristic reactions of Potassium cations K^+

Content	Description	Observation	Explanation
<p>1. Potassium hexanitritocobaltate(III) $Na_3[Co(NO_2)_6]$. The NH_4^+ ions prevent the determination, because ammonia gives similar sediment. <i>Non specific, concentration limit $c = 1:50000$.</i></p>	<p>Add 3-4 drops of reagent to 2-3 drops of potassium salt solution.</p> <p>Analyzed solution should have $pH > 7$ because reagent easily decomposed: $Na_3[Co(NO_2)_6] + 3NaOH = Co(OH)_3 \downarrow + 6NaNO_2$.</p> <p>At $pH < 7$ unstable acid forms which not reacts with K^+ ions: $Na_3[Co(NO_2)_6] \downarrow + 3HCl = H_3[Co(NO_2)_6] \text{ (solution)} + 3NaCl$; $2H_3[Co(NO_2)_6] + 4HCl = 5NO + 2CoCl_2 + 7NO_2 + 5H_2O$.</p>		$2K^+ + Na^+ + [Co(NO_2)_6]^{3-} = K_2Na[Co(NO_2)_6] \downarrow$; $2KCl + Na_3[Co(NO_2)_6] = K_2Na[Co(NO_2)_6] \downarrow + 2NaCl$.
<p>2. Sodium hydrotartrate $NaHC_4H_4O_6$. The NH_4^+ ions prevent the determination, because ammonia gives similar sediment. <i>Non specific, low concentration limit $c = 1:1000$.</i></p>	<p>Add to 4-5 drops of K^+ salt solution the same volume of reagent.</p> <p>Conditions:</p> <ul style="list-style-type: none"> - $pH = 5 \div 7$; - Ammonia ions absence, because the similar sediment forms: $NH_4Cl + NaHC_4H_4O_6 = NH_4HC_4H_4O_6 \downarrow + NaCl$; - Cooling; - Crystallization center formation (rub the tube inner surface by glass stick). 		$K^+ + HC_4H_4O_6^- = KHC_4H_4O_6 \downarrow$; $KCl + NaHC_4H_4O_6 = KHC_4H_4O_6 \downarrow + NaCl$.

1.3.3. Characteristic reactions of Sodium cations Na^+

Content	Description	Observation	Explanation
<p>1. Potassium hexahydroxostibiate $\text{K}[\text{Sb}(\text{OH})_6]$ <i>Non specific</i> - the NH_4^+ ions prevent the determination, because the ammonia salts are hydrolyzed in acidic medium. $\text{K}[\text{Sb}(\text{OH})_6]$ decomposes forming sediment of methastibate acid $[\text{Sb}(\text{OH})_6]^- + \text{H}^+ = \text{HSbO}_3 \downarrow + 3\text{H}_2\text{O}$; pH 5-7. <i>Concentration limit</i> $c=1:3300$.</p>	<p>Add 5-6 drops of NaCl solution to 10-15 drops $\text{K}[\text{Sb}(\text{OH})_6]$; rub the tube well by glass stick; cooling tube content in cold water.</p>		$\text{Na}^+ + [\text{Sb}(\text{OH})_6]^- = \text{Na}[\text{Sb}(\text{OH})_6] \downarrow$ $\text{NaCl} + \text{K}[\text{Sb}(\text{OH})_6] = \text{Na}[\text{Sb}(\text{OH})_6] \downarrow + \text{KCl}.$
<p>2. Zinc-Uranyl-Acetate $\text{Zn}(\text{UO}_2)_3(\text{CH}_3\text{COO})_8$ forms crystalline sediment. Reaction is <i>sensitive</i> (concentration limit $c=1:4000$), <i>highly specific</i> (acetic acid medium).</p>	<p>Initialization of crystallization by rubbing of tube sides by glass stick. Add 8-10 drops of <i>Zinc-Uranyl-Acetate (reagent - in lab fume hood)</i> to 3-4 drops of sodium salt solution and rub of tube sides by glass stick.</p>		$\text{Na}^+ + \text{Zn}^{2+} + 3 \text{UO}_2^{2+} + 8\text{CH}_3\text{COO}^- + \text{CH}_3\text{COOH} + 9\text{H}_2\text{O} = =$ $\text{NaZn}(\text{UO}_2)_3(\text{CH}_3\text{COO})_9 \cdot 9\text{H}_2\text{O} \downarrow;$ $\text{NaCl} + \text{Zn}(\text{UO}_2)_3(\text{CH}_3\text{COO})_8 + \text{CH}_3\text{COOH} + 9\text{H}_2\text{O} =$ $\text{NaZn}(\text{UO}_2)_3(\text{CH}_3\text{COO})_9 \cdot 9\text{H}_2\text{O} \downarrow + \text{HCl}.$

1.4. THE SECOND GROUP OF CATIONS

1.4.1. Action of group reagent $(\text{NH}_4)_2\text{HPO}_4$

Content	Description	Observation	Explanation
<p>The reaction is carried out in presence of mixture $\text{NH}_3 \cdot \text{H}_2\text{O} + \text{NH}_4\text{Cl}$ (buffer solution pH=9-10). At such conditions the cations Ca^{2+}, Sr^{2+}, Fe^{2+}, Fe^{3+}, Al^{3+} are deposited as orthophosphates: $\text{Ca}_3(\text{PO}_4)_2 \downarrow$, $\text{Sr}_3(\text{PO}_4)_2 \downarrow$, $\text{Fe}_3(\text{PO}_4)_2 \downarrow$, $\text{FePO}_4 \downarrow$, $\text{AlPO}_4 \downarrow$, and cation Mg^{2+}, Mn^{2+} are deposited as double salts of Magnesium-ammonium phosphate and Manganese-ammonium phosphate: $\text{MgNH}_4\text{PO}_4 \downarrow$, $\text{MnNH}_4\text{PO}_4 \downarrow$</p>	<p>Take 7 tubes. Add 3-5 drops $\text{NH}_3 \cdot \text{H}_2\text{O}$, 3-5 drops NH_4Cl and 4-5 drops $(\text{NH}_4)_2\text{HPO}_4$. Add 2-3 drops MgCl_2 in the 1st tube, CaCl_2 – in the 2^d one, $\text{Sr}(\text{NO}_3)_2$ – in the 3^d, MnSO_4 – in the 4th, FeSO_4 (bottle with this solution is placed in lab fume hood) – in the 5th, FeCl_3 – in the 6th, $\text{Al}_2(\text{SO}_4)_3$ – in the 7th. Note on the colour of sediments.</p>	<p>1st tube _____ colour of sediment</p> <p>2^d tube _____ colour of sediment</p> <p>3^d tube _____ colour of sediment</p> <p>4th tube _____ colour of sediment</p> <p>5th tube _____ colour of sediment</p> <p>6th tube _____ colour of sediment</p> <p>7th tube _____ colour of sediment</p>	<p>1. $\text{MgCl}_2 + (\text{NH}_4)_2\text{HPO}_4 + \text{NH}_3 \cdot \text{H}_2\text{O} = \text{MgNH}_4\text{PO}_4 \downarrow + 2\text{NH}_4\text{Cl} + \text{H}_2\text{O}$; $\text{Mg}^{2+} + \text{HPO}_4^{2-} + \text{NH}_3 \cdot \text{H}_2\text{O} = \text{MgNH}_4\text{PO}_4 \downarrow + \text{H}_2\text{O}$.</p> <p>2. $3\text{CaCl}_2 + 2(\text{NH}_4)_2\text{HPO}_4 + 2\text{NH}_3 \cdot \text{H}_2\text{O} = \text{Ca}_3(\text{PO}_4)_2 \downarrow + 6\text{NH}_4\text{Cl} + 2\text{H}_2\text{O}$; $3\text{Ca}^{2+} + 2\text{HPO}_4^{2-} + 2\text{NH}_3 \cdot \text{H}_2\text{O} = \text{Ca}_3(\text{PO}_4)_2 \downarrow + 2\text{NH}_4^+ + 2\text{H}_2\text{O}$;</p> <p>3. $3\text{Sr}(\text{NO}_3)_2 + 2(\text{NH}_4)_2\text{HPO}_4 + 2\text{NH}_3 \cdot \text{H}_2\text{O} = \text{Sr}_3(\text{PO}_4)_2 \downarrow + 6\text{NH}_4\text{NO}_3 + 2\text{H}_2\text{O}$; $3\text{Sr}^{2+} + 2\text{HPO}_4^{2-} + 2\text{NH}_3 \cdot \text{H}_2\text{O} = \text{Sr}_3(\text{PO}_4)_2 \downarrow + 2\text{NH}_4^+ + 2\text{H}_2\text{O}$;</p> <p>4. $\text{MnSO}_4 + (\text{NH}_4)_2\text{HPO}_4 + \text{NH}_3 \cdot \text{H}_2\text{O} = \text{MnNH}_4\text{PO}_4 \downarrow + (\text{NH}_4)_2\text{SO}_4 + \text{H}_2\text{O}$; $\text{Mn}^{2+} + \text{HPO}_4^{2-} + \text{NH}_3 \cdot \text{H}_2\text{O} = \text{MnNH}_4\text{PO}_4 \downarrow + \text{H}_2\text{O}$.</p> <p>5. $3\text{FeSO}_4 + 2(\text{NH}_4)_2\text{HPO}_4 + 2\text{NH}_3 \cdot \text{H}_2\text{O} = \text{Fe}_3(\text{PO}_4)_2 \downarrow + 3(\text{NH}_4)_2\text{SO}_4 + 2\text{H}_2\text{O}$; $3\text{Fe}^{2+} + 2\text{HPO}_4^{2-} + 2\text{NH}_3 \cdot \text{H}_2\text{O} = \text{Fe}_3(\text{PO}_4)_2 \downarrow + 2\text{NH}_4^+ + 2\text{H}_2\text{O}$;</p> <p>6. $\text{FeCl}_3 + (\text{NH}_4)_2\text{HPO}_4 + \text{NH}_3 \cdot \text{H}_2\text{O} = \text{FePO}_4 \downarrow + 3\text{NH}_4\text{Cl} + \text{H}_2\text{O}$ $\text{Fe}^{3+} + \text{HPO}_4^{2-} + \text{NH}_3 \cdot \text{H}_2\text{O} = \text{FePO}_4 \downarrow + \text{NH}_4^+ + \text{H}_2\text{O}$;</p> <p>7. $\text{Al}_2(\text{SO}_4)_3 + 2(\text{NH}_4)_2\text{HPO}_4 + 2\text{NH}_3 \cdot \text{H}_2\text{O} = 2\text{AlPO}_4 \downarrow + 3(\text{NH}_4)_2\text{SO}_4 + 3\text{H}_2\text{O}$; $\text{Al}^{3+} + \text{HPO}_4^{2-} + \text{NH}_3 \cdot \text{H}_2\text{O} = \text{AlPO}_4 \downarrow + \text{NH}_4^+ + \text{H}_2\text{O}$.</p>

1.4.2. Characteristic reactions of Magnesium cations Mg^{2+}

Content	Description	Observation	Explanation
<p>Group reagent $(NH_4)_2HPO_4$ at pH 9-10 (buffer mixture $NH_3 \cdot H_2O$ + NH_4Cl) deposits Magnesium ions as crystalline precipitate $MgNH_4PO_4$. That is the qualitative reaction of Mg^{2+} ions is the reaction that is similar to the action of group reagent.</p>	<p>Add 3-5 drops $NH_3 \cdot H_2O$, 3-5 drops NH_4Cl and 2-3 drops $MgCl_2$.</p>		$MgCl_2 + (NH_4)_2HPO_4 + NH_3 \cdot H_2O = MgNH_4PO_4 \downarrow + 2NH_4Cl + H_2O;$ $Mg^{2+} + HPO_4^{2-} + NH_3 \cdot H_2O = MgNH_4PO_4 \downarrow + H_2O.$

1.4.3. Characteristic reactions of Calcium cations Ca^{2+}

Content	Description	Observation	Explanation
<p>1. Sulfate acid H_2SO_4 or soluble sulfates in presence of acetone give $CaSO_4$ white precipitate. It is soluble in concentrated ammonium sulfate $(NH_4)_2SO_4$ solution in the result of complex salt $(NH_4)_2[Ca(SO_4)_2]$ forming. <i>Reaction is selective; the same effect is observed of Ca^{2+} cations.</i></p>	<p>Add 2-3 drops of Sodium sulfate Na_2SO_4 to 4-5 drops of Calcium salt solution.</p> <p>Add 10-15 drops of saturated $(NH_4)_2SO_4$ (bottle is placed in lab fume hood) in a tube, boil in a water bath 1 min, and add to a tube with sedimentation $CaSO_4$.</p>		$CaCl_2 + Na_2SO_4 = CaSO_4 \downarrow + 2NaCl.$ $Ca^{2+} + SO_4^{2-} = CaSO_4 \downarrow$ $CaSO_4 \downarrow + (NH_4)_2SO_4 = (NH_4)_2[Ca(SO_4)_2]$ $CaSO_4 \downarrow + 2NH_4^+ + SO_4^{2-} = 2NH_4^+ + [Ca(SO_4)_2]^{2-}.$

<p>2. Ammonium oxalate $(\text{NH}_4)_2\text{C}_2\text{O}_4$ gives white crystalline precipitate with Calcium ions. It is soluble in mineral acids but is insoluble in acetic acid. <i>Reaction is selective the same effect is observed of Sr^{2+} cations.</i></p>	<p>Add 2-3 drops of Ammonium oxalate to 3-4 drops of CaCl_2 solution. Dissolve obtained sediment, adding by drops HCl.</p>		<p>$\text{CaCl}_2 + (\text{NH}_4)_2\text{C}_2\text{O}_4 = \text{CaC}_2\text{O}_4\downarrow + 2\text{NH}_4\text{Cl};$ $\text{Ca}^{2+} + \text{C}_2\text{O}_4^{2-} = \text{CaC}_2\text{O}_4\downarrow.$</p> <p>$\text{CaC}_2\text{O}_4\downarrow + 2\text{HCl} = \text{CaCl}_2 + \text{H}_2\text{C}_2\text{O}_4;$ $\text{CaC}_2\text{O}_4\downarrow + 2\text{H}^+ = \text{Ca}^{2+} + \text{H}_2\text{C}_2\text{O}_4.$</p>
--	--	--	---

1.4.4. Characteristic reactions of Strontium cations Sr^{2+}

Content	Description	Observation	Explanation
<p>1. Sulfate acid H_2SO_4 and soluble sulfates give SrSO_4 precipitate. It is insoluble in $(\text{NH}_4)_2\text{SO}_4$ concentrated solution. <i>Reaction is selective; the same effect is observed of Ca^{2+} cations.</i></p>	<p>Add 2-3 drops of Sodium sulfate Na_2SO_4 to 4-5 drops of $\text{Sr}(\text{NO}_3)_2$ solution.</p>		<p>$\text{Sr}(\text{NO}_3)_2 + \text{H}_2\text{SO}_4 = \text{SrSO}_4\downarrow + 2\text{HNO}_3;$ $\text{Sr}^{2+} + \text{SO}_4^{2-} = \text{SrSO}_4\downarrow.$</p>
<p>2. Ammonium oxalate $(\text{NH}_4)_2\text{C}_2\text{O}_4$ gives crystalline precipitate with Sr^{2+} ions. It is soluble in mineral acids but insoluble in cold acetic acid. <i>Reaction is selective; the same effect is observed of Ca^{2+} cations.</i></p>	<p>Add 2-3 drops of Ammonium oxalate to 3-4 drops of $\text{Sr}(\text{NO}_3)_2$ solution. Add by drops HCl till dissolving of sediment.</p>		<p>$\text{Sr}(\text{NO}_3)_2 + (\text{NH}_4)_2\text{C}_2\text{O}_4 = \text{SrC}_2\text{O}_4\downarrow + 2\text{NH}_4\text{NO}_3;$ $\text{Sr}^{2+} + \text{C}_2\text{O}_4^{2-} = \text{SrC}_2\text{O}_4\downarrow;$</p> <p>$\text{SrC}_2\text{O}_4\downarrow + 2\text{HCl} = \text{SrCl}_2 + \text{H}_2\text{C}_2\text{O}_4;$ $\text{SrC}_2\text{O}_4\downarrow + 2\text{H}^+ = \text{H}_2\text{C}_2\text{O}_4 + \text{Sr}^{2+}.$</p>
<p>3. Gypsum Water (saturated water solution of CaSO_4) gives</p>	<p>Add 1-2 mL of Gypsum water (reagent - in lab)</p>		<p>$\text{Sr}(\text{NO}_3)_2 + \text{CaSO}_4 = \text{SrSO}_4\downarrow + 2\text{Ca}(\text{NO}_3)_2;$</p>

sediment SrSO_4 in form of white turbidity. Gypsum CaSO_4 is a slightly soluble substance ($\text{SP}=6,1 \cdot 10^{-5}$), but more soluble, that SrSO_4 ($\text{SP}=2,8 \cdot 10^{-7}$). Therefore concentration of SO_4^{2-} in CaSO_4 saturated solution is sufficient for exceeding of SrSO_4 solubility product (SP). <i>Reaction is selective, because Ag^+, Pb^{2+} gives insoluble sulfates too.</i>	fume hood) to 5-10 drops of Strontium salt solution. Heat mixture in a water bath 3-5 min.		$\text{Sr}^{2+} + \text{SO}_4^{2-} = \text{SrSO}_4 \downarrow$.
--	--	--	--

1.4.5. Characteristic reactions of Manganese cations Mn^{2+}

Content	Description	Observation	Explanation				
1. Sodium Bismuthate NaBiO_3 oxidizes Mn^{2+} to MnO_4^- in diluted nitrate acid medium. <i>Reaction is very sensitive and high specific.</i>	Add 15-20 drops of diluted Nitrate acid, 1-2 drops of MnSO_4 and a few grains of solid Sodium Bismuthate (volume as one match-head) (reagent - in lab fume hood).		$2\text{Mn}(\text{NO}_3)_2 + 5\text{NaBiO}_3 + 16\text{HNO}_3 = 2\text{HMnO}_4 + 5\text{Bi}(\text{NO}_3)_3 + 5\text{NaNO}_3 + 7\text{H}_2\text{O};$ $2\text{Mn}^{2+} + 5\text{BiO}_3^- + 14\text{H}^+ = 2\text{MnO}_4^- + 5\text{Bi}^{3+} + 7\text{H}_2\text{O};$ <table style="border-collapse: collapse; margin-left: 20px;"> <tr> <td style="border-right: 1px solid black; padding-right: 5px;">$\text{Mn}^{2+} - 5\text{e}^- \rightarrow \text{Mn}^{7+}$</td> <td style="padding-left: 5px; text-align: center;">2</td> </tr> <tr> <td style="border-right: 1px solid black; padding-right: 5px;">$\text{Bi}^{5+} + 2\text{e}^- \rightarrow \text{Bi}^{3+}$</td> <td style="padding-left: 5px; text-align: center;">5</td> </tr> </table>	$\text{Mn}^{2+} - 5\text{e}^- \rightarrow \text{Mn}^{7+}$	2	$\text{Bi}^{5+} + 2\text{e}^- \rightarrow \text{Bi}^{3+}$	5
$\text{Mn}^{2+} - 5\text{e}^- \rightarrow \text{Mn}^{7+}$	2						
$\text{Bi}^{5+} + 2\text{e}^- \rightarrow \text{Bi}^{3+}$	5						
2. Hydrogen peroxide H_2O_2 oxidizes Mn^{2+} up to Mn^{4+} in ammonia or alkaline medium and sedimentates Manganese in the form mixed oxide-hydroxide. The	lab fume hood) to 2-3		$\text{Mn}(\text{NO}_3)_2 + \text{H}_2\text{O}_2 + 2 \text{NH}_3 \cdot \text{H}_2\text{O} = \text{H}_2\text{MnO}_3 \downarrow + 2\text{NH}_4\text{NO}_3 + \text{H}_2\text{O};$ $\text{Mn}^{2+} + \text{H}_2\text{O}_2 + 2 \text{NH}_3 \cdot \text{H}_2\text{O} = \text{H}_2\text{MnO}_3 \downarrow + 2\text{NH}_4^+ + \text{H}_2\text{O}.$				

black precipitate is formed. <i>Reaction is selective; the same effect is observed of cations formed insoluble hydroxides with alkali (Fe(OH)₃, Cu(OH)₂ etc.).</i>	drops of MnSO ₄ solution. Heat mixture in a water bath 1-2 min.		$\text{Mn}^{2+} - 2e \rightarrow \text{Mn}^{4+} \quad \quad 1$ $2\text{O}^{1-} + 2e \rightarrow 2\text{O}^{2-} \quad \quad 1$
---	---	--	---

1.4.6. Characteristic reactions of Ferrous cations Fe²⁺

Content	Description	Observation	Explanation
1. Alkalis (NaOH, KOH) deposits Fe ²⁺ ions. This reagent is used for separation of Ferrous ions. <i>Reaction is selective; the same effect is observed of cations formed insoluble hydroxides in alkaline medium (Ni(OH)₂, Mn(OH)₂ etc.).</i>	Add 3-5 drops of NaOH, 1-2 drops of FeSO ₄ (reagent - in lab fume hood). Observe the colour changing as the result of oxidation of Fe ²⁺ to Fe ³⁺ by air oxygen.		$\text{FeSO}_4 + 2\text{NaOH} = \text{Fe(OH)}_2\downarrow + \text{Na}_2\text{SO}_4;$ $\text{Fe}^{2+} + 2\text{OH}^- = \text{Fe(OH)}_2\downarrow.$ Oxidation of sediment in the air: $4\text{Fe(OH)}_2 + \text{O}_2 + 2 \text{H}_2\text{O} = 4\text{Fe(OH)}_3\downarrow.$ $\text{Fe}^{2+} - 1e \rightarrow \text{Fe}^{3+} \quad \quad 4$ $\text{O}_2 + 4e \rightarrow 2\text{O}^{2-} \quad \quad 1$
2. Potassium ferrocyanide K₃[Fe(CN)₆] (red blood salt) gives precipitate Fe ₃ [Fe(CN) ₆] ₂ (<i>Turnbull's blue</i>) in slightly acidic medium). It is not soluble in acids, but decomposes in alkalis forming Fe(OH) ₂ precipitate. <i>Reaction is high sensitive and specific.</i>	Add 1 drop of HCl and 2-3 drops of Potassium ferrocyanide to 2-3 drops of FeSO ₄ (reagent - in lab fume hood).		$3\text{FeSO}_4 + 2\text{K}_3[\text{Fe(CN)}_6] = \text{Fe}_3[\text{Fe(CN)}_6]_2\downarrow + 3\text{K}_2\text{SO}_4;$ $3\text{Fe}^{2+} + 2[\text{Fe(CN)}_6]^{3-} = \text{Fe}_3[\text{Fe(CN)}_6]_2\downarrow.$

1.4.7. Characteristic reactions of Ferric cations Fe^{3+}

Content	Description	Observation	Explanation
<p>1. Alkalis (NaOH, KOH) deposits Fe^{3+} ions. This reagent is used for separation of Ferric ions. <i>Reaction is selective; the same effect is observed of cations formed insoluble hydroxides in alkaline medium ($\text{Co}(\text{OH})_2$, $\text{Mg}(\text{OH})_2$ etc.).</i></p>	Add 3-5 drops of NaOH to 1-2 drops of FeCl_3 .		$\text{FeCl}_3 + 3\text{NaOH} = \text{Fe}(\text{OH})_3\downarrow + 3\text{NaCl};$ $\text{Fe}^{3+} + 3\text{OH}^- = \text{Fe}(\text{OH})_3\downarrow.$
<p>2. Potassium ferricyanide $\text{K}_4[\text{Fe}(\text{CN})_6]$ (yellow blood salt) gives precipitate (<i>Prussian blue</i>) with Fe^{3+} ions in acid medium. It isn't soluble in acids but is decomposed in alkalis. <i>Reaction is highly specific and very sensitive (minimum concentration $c = 1:1000000$).</i></p>	Add 1 drop of HCl and 2-3 drops of potassium ferricyanide to 2-3 drops of FeCl_3 solution.		$4\text{FeCl}_3 + 3\text{K}_4[\text{Fe}(\text{CN})_6] = \text{Fe}_4[\text{Fe}(\text{CN})_6]_3\downarrow + 12\text{KCl};$ $4\text{Fe}^{3+} + 3[\text{Fe}(\text{CN})_6]^{4-} = \text{Fe}_4[\text{Fe}(\text{CN})_6]_3\downarrow.$
<p>3. Ammonium or Potassium Thiocyanide (NH_4SCN or KSCN) gives the solution of blood colour. <i>Reaction is very sensitive and specific.</i></p>	Add 1 drop of HCl solution and 2-3 drops NH_4SCN to 2-3 drops of FeCl_3 .		$\text{FeCl}_3 + 6\text{NH}_4\text{SCN} = (\text{NH}_4)_3[\text{Fe}(\text{SCN})_6] + 3\text{NH}_4\text{Cl};$ $\text{Fe}^{3+} + 6\text{SCN}^- = [\text{Fe}(\text{SCN})_6]^{3-}.$

1.4.8. Characteristic reactions of Aluminium cations Al^{3+}

Content	Description	Observation	Explanation
<p>1. Alkalis (NaOH, KOH) form gelatinous precipitate $\text{Al}(\text{OH})_3$. Added excess of alkali, sediment dissolved due to amphoteric character of aluminium compounds. <i>Reaction is selective; the same effect is observed of cations formed amphoteric hydroxides, soluble in alkali ($\text{Zn}(\text{OH})_2$, $\text{Pb}(\text{OH})_2$).</i></p>	<p>Add solution of $\text{Al}_2(\text{SO}_4)_3$ (by drops) to 1-2 drops of alkali as long as white translucent precipitate is formed.</p> <p>Add by drops alkaline solution until sediment dissolves completely.</p>		$\text{Al}_2(\text{SO}_4)_3 + 6\text{NaOH} = 2\text{Al}(\text{OH})_3\downarrow + 3\text{Na}_2\text{SO}_4;$ $\text{Al}^{3+} + 3\text{OH}^- = \text{Al}(\text{OH})_3\downarrow$ <p>In excess of alkali:</p> $\text{Al}(\text{OH})_3\downarrow + \text{NaOH} = \text{Na}[\text{Al}(\text{OH})_4];$ $\text{Al}(\text{OH})_3\downarrow + \text{OH}^- = [\text{Al}(\text{OH})_4]^-.$
<p>2. Ammonium chloride NH_4Cl. Aluminates hydrolyze in water solutions. Equilibrium is shifted to the left or to the right depends on concentrations of ions H^+ and OH^- in solution, i.e. pH of solution. If solid NH_4Cl add to alkaline solution of Aluminate, NH_4Cl reacts with NaOH firstly. But formation of $\text{NH}_3 \cdot \text{H}_2\text{O}$ leads to pH reducing to 7-8 and equilibrium shifts to $\text{Al}(\text{OH})_3$. The best results are observed at heating. <i>Reaction is selective, interferences cations given insoluble hydroxides.</i></p>	<p>Add $\text{Al}_2(\text{SO}_4)_3$ solution by drops to 1-2 drops of alkali until $\text{Al}(\text{OH})_3$ precipitate.</p> <p>Add by drops alkaline solution until sediment dissolved completely.</p> <p>Add 2-3 mL of DW and NH_4Cl (solid) to the saturation (a few crystals fall down and don't solute completely). Heat the mixture in a water bath.</p>		<p>The 1st step – sedimentation of $\text{Al}(\text{OH})_3$:</p> $\text{Al}_2(\text{SO}_4)_3 + 6\text{NaOH} = 2\text{Al}(\text{OH})_3\downarrow + 3\text{Na}_2\text{SO}_4;$ $\text{Al}^{3+} + 3\text{OH}^- = \text{Al}(\text{OH})_3\downarrow$ <p>Dissolving in excess of alkali:</p> $\text{Al}(\text{OH})_3\downarrow + \text{NaOH} = \text{Na}[\text{Al}(\text{OH})_4];$ $\text{Al}(\text{OH})_3\downarrow + \text{OH}^- = [\text{Al}(\text{OH})_4]^-;$ <p>Sedimenting of $\text{Al}(\text{OH})_3$ in saturated NH_4Cl solution:</p> $\text{Na}[\text{Al}(\text{OH})_4] + \text{NH}_4\text{Cl} = \text{Al}(\text{OH})_3\downarrow + \text{NH}_3 \cdot \text{H}_2\text{O} + \text{NaCl};$ $[\text{Al}(\text{OH})_4]^- + \text{NH}_4^+ = \text{Al}(\text{OH})_3\downarrow + \text{NH}_3 \cdot \text{H}_2\text{O}.$

1.5. THE THIRD GROUP OF CATIONS

1.5.1. Action of group reagent $(\text{NH}_4)_2\text{HPO}_4$

Content	Description	Observation	Explanation				
<p>The cations Cu^{2+}, Zn^{2+}, Co^{2+}, Ni^{2+} are precipitated by $(\text{NH}_4)_2\text{HPO}_4$ as orthophosphates $\text{Cu}_3(\text{PO}_4)_2$, $\text{Zn}_3(\text{PO}_4)_2$, $\text{Co}_3(\text{PO}_4)_2$, and $\text{Ni}_3(\text{PO}_4)_2$.</p> <p>$\text{Co}_3(\text{PO}_4)_2$ and $\text{Ni}_3(\text{PO}_4)_2$ are unstable in open air in the presence of oxygen, changing colour in the result of cations oxidation.</p>	<p>Take 4 tubes. Add 1-2 drops $\text{NH}_3 \cdot \text{H}_2\text{O}$, 1-2 drops NH_4Cl, and 4-5 drops $(\text{NH}_4)_2\text{HPO}_4$.</p> <p>Add 2-3 drops CuSO_4 in the 1st tube, ZnSO_4 – in the 2^d one, CoCl_2 – in the 3^d, and NiSO_4 – in the 4th. Note on the colour of sediments and its stability.</p>	<p>1st tube _____ colour of sediment</p> <p>2^d tube _____ colour of sediment</p> <p>3^d tube _____ colour of sediment</p> <p>_____</p> <p>Stability of colour</p> <p>4th tube _____ colour of sediment</p> <p>_____</p> <p>Stability of colour</p>	<p>1. $3\text{MeSO}_4 + 2(\text{NH}_4)_2\text{HPO}_4 + 2\text{NH}_3 \cdot \text{H}_2\text{O} = \text{Me}_3(\text{PO}_4)_2\downarrow + 3(\text{NH}_4)_2\text{SO}_4 + 2\text{H}_2\text{O}$; $3\text{Me}^{2+} + 2\text{HPO}_4^{2-} + 2\text{NH}_3 \cdot \text{H}_2\text{O} = \text{Me}_3(\text{PO}_4)_2\downarrow + 2\text{NH}_4^+ + 2\text{H}_2\text{O}$ (where Me – Cu^{2+}; Zn^{2+}, Ni^{2+});</p> <p>2. $3\text{CoCl}_2 + 2(\text{NH}_4)_2\text{HPO}_4 + 2\text{NH}_3 \cdot \text{H}_2\text{O} = \text{Co}_3(\text{PO}_4)_2\downarrow + 6\text{NH}_4\text{Cl} + 2\text{H}_2\text{O}$; $3\text{Co}^{2+} + 2\text{HPO}_4^{2-} + 2\text{NH}_3 \cdot \text{H}_2\text{O} = \text{Co}_3(\text{PO}_4)_2\downarrow + 2\text{NH}_4^+ + 2\text{H}_2\text{O}$; Oxidation of precipitate in the air: $4\text{Co}_3(\text{PO}_4)_2\downarrow + 3\text{O}_2 + 6\text{H}_2\text{O} = 4(\text{CoOH})_3(\text{PO}_4)_2$.</p> <table style="display: inline-table; vertical-align: middle;"> <tr> <td style="padding-right: 10px;">$3\text{Co}^{2+} - 3\text{e}^- \rightarrow \text{Co}^{3+}$</td> <td style="border-left: 1px solid black; padding-left: 10px; text-align: center;">4</td> </tr> <tr> <td style="padding-right: 10px;">$\text{O}_2^0 + 4\text{e}^- \rightarrow 2\text{O}^{2-}$</td> <td style="border-left: 1px solid black; padding-left: 10px; text-align: center;">3</td> </tr> </table>	$3\text{Co}^{2+} - 3\text{e}^- \rightarrow \text{Co}^{3+}$	4	$\text{O}_2^0 + 4\text{e}^- \rightarrow 2\text{O}^{2-}$	3
$3\text{Co}^{2+} - 3\text{e}^- \rightarrow \text{Co}^{3+}$	4						
$\text{O}_2^0 + 4\text{e}^- \rightarrow 2\text{O}^{2-}$	3						
Solubility of sediments obtained at the action of group reagent							
Content	Description	Observation	Explanation				
<p>All above mentioned precipitates are dissolved in excess of $\text{NH}_3 \cdot \text{H}_2\text{O}$. They are soluble in mineral acids (HCl, H_2SO_4) and acetic acid CH_3COOH.</p>	<p>Add 10-15 drops of water ammonia to content of tubes # 1-4. Describe the visual effects.</p>		<p>1. $\text{Me}_3(\text{PO}_4)_2\downarrow + 12\text{NH}_3 \cdot \text{H}_2\text{O} = [\text{Me}(\text{NH}_3)_4]_3(\text{PO}_4)_2 + 12\text{H}_2\text{O}$ (where Me – Cu^{2+}, Zn^{2+});</p> <p>$\text{Me}_3(\text{PO}_4)_2\downarrow + 12\text{NH}_3 \cdot \text{H}_2\text{O} = 3[\text{Me}(\text{NH}_3)_4]^{2+} + 2\text{PO}_4^{3-} + 12\text{H}_2\text{O}$;</p>				

Zinc orthophosphate due to amphoteric character is soluble in excess of alkali solution.			$2. \text{Me}_3(\text{PO}_4)_2\downarrow + 18 \text{NH}_3\cdot\text{H}_2\text{O} = [\text{Me}(\text{NH}_3)_6]_3(\text{PO}_4)_2 + 18 \text{H}_2\text{O}$ <p>(where Me – Co^{2+}, Ni^{2+});</p> $\text{Me}_3(\text{PO}_4)_2\downarrow + 18 \text{NH}_3\cdot\text{H}_2\text{O} = 3[\text{Me}(\text{NH}_3)_6]^{2+} + 2\text{PO}_4^{3-} + 18 \text{H}_2\text{O};$ $\text{Zn}_3(\text{PO}_4)_2\downarrow + 12\text{NaOH} = 3 \text{Na}_2[\text{Zn}(\text{OH})_4] + 2\text{Na}_3\text{PO}_4;$ $\text{Zn}_3(\text{PO}_4)_2\downarrow + 12\text{OH}^- = 3[\text{Zn}(\text{OH})_4]^{2-} + 2\text{PO}_4^{3-}.$
--	--	--	---

1.5.2. Characteristic reactions of Copper cations Cu^{2+}

Content	Description	Observation	Explanation
<p>1. Water ammonia $\text{NH}_3\cdot\text{H}_2\text{O}$ in stoichiometrical quantity gives precipitate of a basic copper salt; adding in excess – dissolved the sediment. Due to the colour of solution this test is <i>specific reaction of Cu^{2+} and highly sensitive ($c=1:500000$).</i></p>	<p>Add 1-2 drops of water ammonia in test tube and add by drops CuSO_4 solution until precipitate formation. Note on the on its colour.</p> <p>After that add by drops an excess of water ammonia till the complete dissolving of precipitate. What's colour of solution?</p>		<p>The reaction is carried out in two stages:</p> <p>the 1st:</p> $2\text{CuSO}_4 + 2\text{NH}_3\cdot\text{H}_2\text{O} = (\text{CuOH})_2\text{SO}_4\downarrow + (\text{NH}_4)_2\text{SO}_4;$ $2\text{Cu}^{2+} + \text{SO}_4^{2-} + 2\text{NH}_3\cdot\text{H}_2\text{O} = (\text{CuOH})_2\text{SO}_4\downarrow + 2\text{NH}_4^+;$ <p>the 2^d:</p> $(\text{CuOH})_2\text{SO}_4\downarrow + (\text{NH}_4)_2\text{SO}_4 + 6\text{NH}_3\cdot\text{H}_2\text{O} = 2[\text{Cu}(\text{NH}_3)_4]\text{SO}_4 + 8\text{H}_2\text{O};$ $(\text{CuOH})_2\text{SO}_4\downarrow + 2\text{NH}_4^+ + 6\text{NH}_3\cdot\text{H}_2\text{O} = 2[\text{Cu}(\text{NH}_3)_4]^{2+} + 8\text{H}_2\text{O}.$
<p>2. Alkalis (NaOH, KOH) give precipitate with copper(II) ions.</p>	<p>Add 2-3 drops of NaOH to solution of CuSO_4</p>		$\text{CuSO}_4 + 2\text{NaOH} = \text{Cu}(\text{OH})_2\downarrow + \text{Na}_2\text{SO}_4;$

<p>This reaction is <i>selective</i>; uses for separation of Cu^{2+} ions from cations mixture.</p>	<p>until sedimentation. Copper hydroxide has weak amphoteric properties, dissolved in mineral and acetic acids.</p>		$\text{Cu}^{2+} + 2\text{OH}^- = \text{Cu}(\text{OH})_2 \downarrow.$
<p>3. Potassium ferricyanide $\text{K}_4[\text{Fe}(\text{CN})_6]$ (<i>yellow blood salt</i>) forms precipitate with Cu^{2+} ions ($\text{pH} < 7$). Reaction is <i>selective</i>; but highly sensitive ($c=1:2500000$). Precipitate $\text{Cu}_2[\text{Fe}(\text{CN})_6]$ isn't dissolve in diluted acids, but solves in excess of water ammonia.</p>	<p>Add 1 drop of HCl and 2-3 drops of $\text{K}_4[\text{Fe}(\text{CN})_6]$ to 3-4 drops of CuCl_2 solution.</p>		$\begin{aligned} 2\text{CuCl}_2 + \text{K}_4[\text{Fe}(\text{CN})_6] &= \\ \text{Cu}_2[\text{Fe}(\text{CN})_6] \downarrow + 4\text{KCl}; & \\ 2\text{Cu}^{2+} + [\text{Fe}(\text{CN})_6]^{4-} &= \\ \text{Cu}_2[\text{Fe}(\text{CN})_6] \downarrow. & \end{aligned}$
<p>4. Potassium Iodide KI reacts with copper salt formed white precipitate CuI. But reactive mixture has brown-red colour due to free I_2 isolation. Reaction is <i>selective</i>; but highly sensitive ($c=1:300000$).</p>	<p>Add 2-3 drops of KI solution to 3-4 drops of CuSO_4 solution.</p>		$\begin{aligned} 2\text{CuSO}_4 + 4\text{KI} &= 2\text{CuI} + \text{I}_2 + \\ &2\text{K}_2\text{SO}_4; \\ 2\text{Cu}^{2+} + 4\text{I}^- &\rightarrow 2\text{CuI} \downarrow + \text{I}_2. \\ \hline 2\text{I}^- - 2\text{e} &\rightarrow \text{I}_2 & & 1 \\ \text{Cu}^{2+} + 1\text{e} &\rightarrow \text{Cu}^+ & & 2 \end{aligned}$

1.5.3. Characteristic reactions of Zinc cations Zn^{2+}

Content	Description	Observation	Explanation
<p>1. Water ammonia $NH_3 \cdot H_2O$ in stoichiometrical quantity precipitates zinc hydroxide; adding in excess – dissolves the sediment. This test is <i>selective reaction of Zn^{2+}</i>.</p>	<p>Add 1-2 drops of water ammonia in test tube and add by drops $ZnSO_4$ solution until the sedimentation. After then add by drops an excess of ammonia till the complete dissolving. Note on the colour of solution.</p>		<p>The reaction is carried out in two stages: the 1st: $ZnSO_4 + 2NH_3 \cdot H_2O = Zn(OH)_2 \downarrow + (NH_4)_2SO_4$; $Zn^{2+} + 2NH_3 \cdot H_2O = Zn(OH)_2 \downarrow + 2NH_4^{2+}$; the 2^d: $Zn(OH)_2 \downarrow + 4 NH_3 \cdot H_2O \text{ (in excess)} = [Zn(NH_3)_4](OH)_2 + 4H_2O$; $Zn(OH)_2 \downarrow + 4 NH_3 \cdot H_2O = [Zn(NH_3)_4]^{2+} + 4H_2O$.</p>
<p>2. Alkalis ($NaOH$, KOH) give white $Zn(OH)_2$ gelatinous precipitate with Zn^{2+} ions. This sediment has amphoteric properties and soluble in mineral acids and excess of alkalis. This test is <i>selective, used for the separation of Zn^{2+} from cations mixture</i>.</p>	<p>Add slowly 10-12 drops of $ZnSO_4$ solution to 2-3 drops of $NaOH$ or KOH solution up to forming of white precipitate $Zn(OH)_2$. After then add excess of alkali to complete dissolving of sediment obtained before.</p>		<p>The reaction is carried out in two stages: the 1st: $ZnSO_4 + 2NaOH = Zn(OH)_2 \downarrow + Na_2SO_4$; $Zn^{2+} + 2OH^- = Zn(OH)_2 \downarrow$; the 2^d: adding excess of alkali: $Zn(OH)_2 \downarrow + 2NaOH = Na_2[Zn(OH)_4]$; $Zn(OH)_2 \downarrow + 2OH^- = [Zn(OH)_4]^{2-}$.</p>

<p>3. Potassium ferrocyanide $K_3[Fe(CN)_6]$ (red blood salt) forms precipitate of mustard colour with Zn ions (pH<7). It solves in alkalis. This test is selective, Fe^{3+} interferences.</p>	<p>Add 1 drops HCl and 2-3 drops of $K_4[Fe(CN)_6]$ to 2-3 drops of $ZnSO_4$ solution.</p>		$3ZnSO_4 + 2K_3[Fe(CN)_6] = Zn_3[Fe(CN)_6]_2\downarrow + 3K_2SO_4;$ $3Zn^{2+} + 2[Fe(CN)_6]^{3-} = Zn_3[Fe(CN)_6]_2\downarrow.$
--	--	--	---

1.5.4. Characteristic reactions of Cobalt cations Co^{2+}

Content	Description	Observation	Explanation
<p>1. Water ammonia $NH_3 \cdot H_2O$ in stoichiometrical quantity precipitates basic cobalt salt; adding in excess – dissolves the sediment. Test is <i>selective</i>.</p>	<p>Add 1-2 drops of water ammonia in test tube and add by drops $CoCl_2$ solution till the sedimenting of precipitate. After then add by drops an excess of ammonia till the complete dissolving. Note on the colour of solution, which is changing quickly in the result of oxidation $Co^{2+} \rightarrow Co^{3+}$.</p>		<p>The reaction is carried out in two stages: the 1st: $CoCl_2 + NH_3 \cdot H_2O = CoOHCl\downarrow + NH_4Cl;$ $Co^{2+} + Cl^- + NH_3 \cdot H_2O = CoOHCl\downarrow + NH_4^+.$ the 2^d: $CoOHCl\downarrow + 7 NH_3 \cdot H_2O = [Co(NH_3)_6](OH)_2 + NH_4Cl + 6H_2O;$ $CoOHCl\downarrow + 7 NH_3 \cdot H_2O = [Co(NH_3)_6]^{2+} + 2OH^- + NH_4^+ + Cl^-.$</p>
<p>2. Alkalis ($NaOH$, KOH) give gelatinous precipitate of basic salt with Co^{2+} ions. Excess of alkali converts $CoOHCl$ at heating in $Co(OH)_2$ of pink</p>	<p>Add 1 drop of $NaOH$ to 3-4 drops of $CoCl_2$ solution. Note on the colour of sediment. Add excess of alkali. Observe change of</p>		$CoCl_2 + NaOH = CoOHCl\downarrow + NaCl;$ $Co^{2+} + OH^- + Cl^- = CoOHCl\downarrow.$ $CoOHCl\downarrow + NaOH = Co(OH)_2\downarrow + NaCl;$

colour. Test is <i>selective</i> .	sediment colour. Oxidation by air oxygen, Co(OH)_2 converts to Co(OH)_3 . Note on the colour of solution.		$\text{CoOHCl} \downarrow + \text{OH}^- = \text{Co(OH)}_2 \downarrow + \text{Cl}^-$ Oxidation: $4\text{Co(OH)}_2 + \text{O}_2 + 2\text{H}_2\text{O} = 4\text{Co(OH)}_3 \downarrow$: $\begin{array}{l l} \text{Co}^{2+} - 1\text{e} \rightarrow \text{Co}^{3+} & 4 \\ \text{O}_2^0 + 4\text{e} \rightarrow 2\text{O}^{2-} & 1 \end{array}$
3. Ammonium or Potassium Thiocyanide (NH_4SCN or KSCN) (<i>Vogel reaction</i>) solution in organic solvents (acetone) gives the dark blue solution. Reaction is <i>very sensitive, but selective</i> . Interference of Fe^{3+} ions can be eliminated by adding some solid potassium fluoride crystals NaF . The colourless hexafluoroferrate(III) $[\text{FeF}_6]^{3-}$ complex does not interfere, because it is more stable than $[\text{Fe(SCN)}_6]^{3-}$. Concentration limit 1 : 100000.	Add 4-5 drops of NH_4SCN saturated solution in acetone (reagent - in lab fume hood) to 3-4 drops of CoCl_2 . Shake up content of tube and wait a few seconds. Note on the colour of upper layer.		$\text{CoCl}_2 + 6\text{NH}_4\text{SCN} = (\text{NH}_4)_4[\text{Co(SCN)}_6] + 2\text{NH}_4\text{Cl};$ $\text{Co}^{2+} + 6\text{SCN}^- = [\text{Co(SCN)}_6]^{4-}.$
4. Potassium Nitrite KNO_2 in presence of acetic acid oxidizes Co^{2+} to Co^{3+} forming with excess of KNO_2 sediment $\text{K}_3[\text{Co(NO}_2)_6]$. Reaction is <i>very sensitive, but not specific</i> .	To 5-7 drops of CoCl_2 solution add solid crystals KNO_2 to saturation. Add concentrated acetic acid (reagent - in lab fume hood) by drops up to formation of precipitate.		$\text{CoCl}_2 + 7\text{KNO}_2 + 2\text{CH}_3\text{COOH} = \text{K}_3[\text{Co(NO}_2)_6] \downarrow + 2\text{KCl} + 2\text{CH}_3\text{COOK} + \text{NO} + \text{H}_2\text{O}.$ $\begin{array}{l l} \text{Co}^{2+} - 1\text{e} \rightarrow \text{Co}^{3+} & 1 \\ \text{N}^{3+} + 1\text{e} \rightarrow \text{N}^{2+} & 1 \end{array}$ $\text{Co}^{2+} + 7\text{NO}_2^- + 3\text{K}^+ + 2\text{CH}_3\text{COOH} = \text{K}_3[\text{Co(NO}_2)_6] \downarrow + 2\text{CH}_3\text{COO}^- + \text{H}_2\text{O} + \text{NO}.$

1.5.5. Characteristic reactions of Nickel cations Ni²⁺

Content	Description	Observation	Explanation
<p>1. Water ammonia NH₃·H₂O in stoichiometrical quantity precipitates basic nickel salt; adding in excess – dissolved the sediment. Test is <i>selective</i>, Cu²⁺ and Co²⁺ ions interfere. Reaction <i>is selective</i>.</p>	<p>Add by drops 2NH₃·H₂O solution to 3-4 drops of NiSO₄ up to basic salt precipitate and then the excess of ammonia up to the dissolving of precipitate.</p>		<p>The reaction is carried out in two stages: the 1st: $2\text{NiSO}_4 + 2\text{NH}_3 \cdot \text{H}_2\text{O} = (\text{NiOH})_2\text{SO}_4 \downarrow + (\text{NH}_4)_2\text{SO}_4$; $2\text{Ni}^{2+} + 2\text{SO}_4^{2-} + 2\text{NH}_3 \cdot \text{H}_2\text{O} = (\text{NiOH})_2\text{SO}_4 \downarrow + 2\text{NH}_4^+$; the 2^d: $(\text{NiOH})_2\text{SO}_4 \downarrow + (\text{NH}_4)_2\text{SO}_4 + 2\text{NH}_3 \cdot \text{H}_2\text{O} = 2[\text{Ni}(\text{NH}_3)_6]\text{SO}_4 + 12\text{H}_2\text{O}$; $(\text{NiOH})_2\text{SO}_4 \downarrow + 2\text{NH}_4^+ + 10\text{NH}_3 \cdot \text{H}_2\text{O} = 2[\text{Ni}(\text{NH}_3)_6]^{2+} + \text{SO}_4^{2-} + 12\text{H}_2\text{O}$.</p>
<p>2. Alkalis (NaOH, KOH) give gelatinous precipitate Ni(OH)₂. Nickel(II) hydroxide dissolves in mineral acids, excess of water ammonia, solutions of ammonia salts, but isn't soluble in excess of alkali. Reaction <i>is selective</i>.</p>	<p>Add by drops NiSO₄ to 1-2 drops of NaOH.</p>		<p>$\text{NiSO}_4 + 2\text{NaOH} = \text{Ni}(\text{OH})_2 \downarrow + \text{Na}_2\text{SO}_4$; $\text{Ni}^{2+} + 2\text{OH}^- = \text{Ni}(\text{OH})_2 \downarrow$.</p>
<p>3. Dimethylglyoxime (Chugaev's reagent) C₄H₈N₂O₂ in ammonia medium (pH=9) forms precipitate of intra-complex salt with Ni²⁺</p>	<p>Add 5-7 drops of water ammonia solution NH₃·H₂O and 3-5 alcohol solution of Chugaev's</p>		<p>$\text{NiSO}_4 + 2\text{C}_4\text{H}_8\text{N}_2\text{O}_2 + 2\text{NH}_3 \cdot \text{H}_2\text{O} = [\text{Ni}(\text{C}_4\text{H}_7\text{N}_2\text{O}_2)_2] \downarrow + (\text{NH}_4)_2\text{SO}_4 + 2\text{H}_2\text{O}$; $\text{Ni}^{2+} + 2\text{C}_4\text{H}_8\text{N}_2\text{O}_2 + 2\text{NH}_3 \cdot \text{H}_2\text{O} = [\text{Ni}(\text{C}_4\text{H}_7\text{N}_2\text{O}_2)_2] \downarrow + 2\text{NH}_4^+ + 2\text{H}_2\text{O}$.</p>

ions. It solves in acids, alkalis, but doesn't dissolve in diluted ammonia solution. Reaction is very sensitive, concentration limit $c=1:300000$. Ions of Fe^{2+} , Fe^{3+} , Cu^{2+} interference of Ni^{2+} determination.	reagent (or Dimethylglyoxime - in lab fume hood) to 2-4 drops of $NiSO_4$ solution. Reaction may be realized in the form of drop test on paper filter.		
--	--	--	--

1.6. THE FOURTH GROUP OF CATIONS

1.6.1. Action of group reagent HCl

Content	Description	Observation	Explanation
The diluted HCl and soluble chlorides deposits Ag^+ and Pb^{2+} in form of white precipitates of $AgCl$ and $PbCl_2$, not soluble in diluted HNO_3 and H_2SO_4 .	Take 2 tubes. Add 2-3 drops of $AgNO_3$ (reagent - in lab fume hood) in the first tube; 2-3 drops $Pb(NO_3)_2$ in the second tube. Add the same quantity of diluted HCl. Formed precipitates of $AgCl$ and $PbCl_2$ divide into 4 tubes each. Study their solubility in hot	1 st tube _____ colour of sediment 2 ^d tube _____ colour of sediment 3 ^d tube _____ _____ 4 th tube _____ _____	$AgNO_3 + HCl = AgCl\downarrow + HNO_3$; $Ag^+ + Cl^- = AgCl\downarrow$; $Pb(NO_3)_2 + 2HCl = PbCl_2\downarrow + 2HNO_3$; $Pb^{2+} + 2Cl^- = PbCl_2\downarrow$; $PbCl_2\downarrow \xrightarrow{t^{\circ}C} Pb^{2+} + 2Cl^-$ (dissolving in boiling water); $AgCl\downarrow + 2NH_3 \cdot H_2O = [Ag(NH_3)_2]Cl + 2H_2O$ (dissolving in concentrated ammonia); $AgCl\downarrow + 2NH_3 \cdot H_2O = [Ag(NH_3)_2]^{2+} + Cl^- + 2H_2O$; $AgCl\downarrow + (NH_4)_2CO_3 = [Ag(NH_3)_2]Cl$

	H ₂ O, concentrated NH ₃ ·H ₂ O (reagent - in lab fume hood), concentrated (NH ₄) ₂ CO ₃ (reagent - in lab fume hood), concentrated HCl (reagent - in lab fume hood). PbCl ₂ in contrast to AgCl dissolves well in hot DW. This technique is used for the separation of AgCl and PbCl ₂ precipitates.	5 th tube _____ _____	$+2\text{H}_2\text{O} + \text{CO}_2\uparrow$ (dissolving in concentrated ammonia carbonate); $\text{AgCl}\downarrow + 2\text{NH}_4^+ + \text{CO}_3^{2-} = [\text{Ag}(\text{NH}_3)_2]^+ + \text{Cl}^- + 2\text{H}_2\text{O} + \text{CO}_2\uparrow$; $\text{AgCl}\downarrow + \text{HCl} = \text{H}[\text{AgCl}_2]$; $\text{AgCl}\downarrow + \text{H}^+ + \text{Cl}^- = \text{H}^+ + [\text{AgCl}_2]^-$; $\text{PbCl}_2\downarrow + 2\text{HCl} = \text{H}_2[\text{PbCl}_4]$; $\text{PbCl}_2\downarrow + 2\text{H}^+ + 2\text{Cl}^- = 2\text{H}^+ + [\text{PbCl}_4]^{2-}$.
	6 th tube _____ _____		
	7 th tube _____ _____		
	8 th tube _____ _____		

1.6.2. Characteristic reactions of Silver cations Ag⁺

Content	Description	Observation	Explanation
1. Water ammonia NH₃·H₂O precipitates silver cation in form of Ag ₂ O, which is dissolved in excess of reactant. Reaction is <i>selective</i> .	Add by drops water ammonia solution to 1-2 drops of AgNO ₃ solution till precipitate formation. Observe dissolving of turbidity at adding of excess of NH ₃ ·H ₂ O.		$\text{AgNO}_3 + \text{NH}_3 \cdot \text{H}_2\text{O} = \text{AgOH}\downarrow + \text{NH}_4\text{NO}_3$; $\text{Ag}^+ + \text{NH}_3 \cdot \text{H}_2\text{O} = \text{AgOH}\downarrow + \text{NH}_4^+$; $2\text{AgOH}\downarrow = \text{Ag}_2\text{O}\downarrow + \text{H}_2\text{O}$; $\text{Ag}_2\text{O}\downarrow + 4\text{NH}_3 \cdot \text{H}_2\text{O} = 2[\text{Ag}(\text{NH}_3)_2]\text{OH} + 3\text{H}_2\text{O}$; $\text{Ag}_2\text{O}\downarrow + 4\text{NH}_3 \cdot \text{H}_2\text{O} = 2[\text{Ag}(\text{NH}_3)_2]^+ + 2\text{OH}^- + 3\text{H}_2\text{O}$.

1.6.3. Characteristic reactions of Lead cations Pb^{2+}

Content	Description	Observation	Explanation
<p>1. Potassium Iodide KI isolates sediment PbI_2. The compound is colourless when dissolved in hot water, but crystallizes on cooling as thin but visibly larger bright yellow flakes, that settle slowly through the liquid — a visual effect often described as "golden rain". This reaction is called sometimes "golden rain". Conditions of re-crystallization: pH less than 7, fast cooling. The test <i>is selective</i>.</p>	<p>Add 1-3 KI drops to 2 drops of $\text{Pb}(\text{NO}_3)_2$, acidified by 1-2 drops of CH_3COOH. Add 5-7 mL DW to precipitate and boil mixture in a water bath. After complete dissolving of precipitate, cool mixture quickly at stream of tap water. For acceleration of re-crystallization rub tube sides of glass stick.</p>		$\text{Pb}(\text{NO}_3)_2 + 2\text{KI} = \text{PbI}_2\downarrow + 2\text{KNO}_3;$ $\text{Pb}^{2+} + 2\text{I}^- = \text{PbI}_2\downarrow.$
<p>2. Potassium Chromate (Dichromate) $K_2\text{CrO}_4$ ($K_2\text{Cr}_2\text{O}_7$) gives yellow precipitate PbCrO_4 with Pb^{2+} ions. This sediment doesn't dissolve in acetic and hydrochloric acids, but solutes in alkali and nitrate acid. Conditions: pH less than 7 (in presence of acetic acid), absence of reducing agents, reduced Cr^{6+} to Cr^{3+} (for example, Fe^{2+}) Reaction <i>is</i></p>	<p>Add 1-2 drops of K_2CrO_4 or $\text{K}_2\text{Cr}_2\text{O}_7$ solution to 1-2 drops of $\text{Pb}(\text{NO}_3)_2$ solution.</p>		$\text{Pb}(\text{NO}_3)_2 + \text{K}_2\text{CrO}_4 = \text{PbCrO}_4\downarrow + 2\text{KNO}_3;$ $\text{Pb}^{2+} + \text{CrO}_4^{2-} = \text{PbCrO}_4\downarrow.$

<p><i>selective</i>, in presence of Ag^+ brick-red sediment Ag_2CrO_4 or dark-red $\text{Ag}_2\text{Cr}_2\text{O}_7$ is precipitated.</p>			
<p>3. Diluted Sulfate Acid H_2SO_4 and soluble sulfates precipitate Pb^{2+} cation in the form of crystalline precipitate PbSO_4. Sediment doesn't dissolve in excess of diluted H_2SO_4, but dissolves in alkalis, ammonia acetate, conc. H_2SO_4 and HCl. Conditions: $\text{pH} < 7$, because PbSO_4 dissolves in alkalis, absence of Ca^{2+} and Sr^{2+} ions in examined solution. <i>Reaction is selective.</i></p>	<p>Add 1-2 drops of diluted H_2SO_4 to 2-3 drops of $\text{Pb}(\text{NO}_3)_2$.</p>		<p>$\text{Pb}(\text{NO}_3)_2 + \text{H}_2\text{SO}_4 = \text{PbSO}_4 \downarrow + 2\text{HNO}_3$; $\text{Pb}^{2+} + \text{SO}_4^{2-} = \text{PbSO}_4$.</p>

1.7. REACTIONS OF THE ANIONS








The methods available for the detection of anions are not as systematic as those which have been described in the previous chapter 1 for cations. No really satisfactory scheme has yet been proposed which permits of the separation of the common anions into major groups, and the subsequent unequivocal separation of each group into its independent constituents. It must, however, be mentioned that it is possible to separate the anions into major groups dependent upon the solubility of their silver salts, or their barium salts. These however, can only be regarded as useful in giving an indication of the limitations of the method and for the confirmation of the results obtained by the simpler procedures to be described below.

Classification of anions

Group	I	II	III
Anions included	SO_4^{2-} ; SO_3^{2-} ; CO_3^{2-} ; PO_4^{3-} ; SiO_3^{2-} ; BO_2^- (or $\text{B}_4\text{O}_7^{2-}$); MoO_4^{2-}	Cl^- , Br^- , I^-	NO_3^- , NO_2^- , CH_3COO^-
Group reagent	BaCl_2 in neutral or weakly alkaline medium.	AgNO_3 in the presence of diluted HNO_3 .	No group reagent, practically all salts are soluble in water.
Properties of precipitates	Barium salts are insoluble in H_2O but soluble in diluted acids (excepting BaSO_4).	Silver salts are insoluble in H_2O and HNO_3 .	Barium and silver salts dissolve in water.


1.7.1. The first group of anions

BaCl_2 as group reagent deposits the first group anions in the form of white precipitates BaSO_4 , BaSO_3 , BaCO_3 , $\text{Ba}_3(\text{PO}_4)_2$, $\text{Ba}(\text{BO}_2)_2$ or BaB_4O_7 , BaSiO_3 , BaMoO_4 . All precipitates are soluble in Hydrochloric HCl and Nitrate HNO_3 acids, excluding BaSO_4 . Precipitates $\text{Ba}_3(\text{PO}_4)_2$, BaCO_3 and $\text{Ba}(\text{BO}_2)_2$ solve in acetic acid CH_3COOH . For example,



Content	Description	Observation	Explanation
Characteristic reactions of Sulfate anions SO_4^{2-}			
1. BaCl_2 (group reagent) in neutral medium deposits SO_4^{2-} in form of precipitate BaSO_4 , insoluble in acids and alkali.	Add 1-2 drops of BaCl_2 solution to 2-3 drops of Na_2SO_4 . Divide the sediment on two test tubes. Add 10-15 HCl drops to the first one, and 5-10 NaOH drops to the second one.		$\text{Na}_2\text{SO}_4 + \text{BaCl}_2 = \text{BaSO}_4\downarrow + 2\text{NaCl}$; $\text{SO}_4^{2-} + \text{Ba}^{2+} = \text{BaSO}_4\downarrow$; $\text{BaSO}_4\downarrow + \text{HCl} \rightarrow \text{no react}$; $\text{BaSO}_4\downarrow + \text{NaOH} \rightarrow \text{no react}$.
2. Soluble Lead salt (i.e., $\text{Pb}(\text{NO}_3)_2$) precipitates SO_4^{2-} anions in the form of PbSO_4 , insoluble in acids, but soluble in alkali (in contrast to BaSO_4).	Add 1-2 drops of $\text{Pb}(\text{NO}_3)_2$ to 2-3 drops of Na_2SO_4 . Add by drops of NaOH until obtained precipitate is dissolved completely.		$\text{Na}_2\text{SO}_4 + \text{Pb}(\text{NO}_3)_2 = \text{PbSO}_4\downarrow + 2\text{NaNO}_3$; $\text{SO}_4^{2-} + \text{Pb}^{2+} = \text{PbSO}_4\downarrow$; $\text{PbSO}_4\downarrow + 4\text{NaOH} (\text{excess}) = \text{Na}_2[\text{Pb}(\text{OH})_4] + \text{Na}_2\text{SO}_4$; $\text{PbSO}_4\downarrow + 4\text{OH}^- = [\text{Pb}(\text{OH})_4]^{2-} + \text{SO}_4^{2-}$.
Characteristic reactions of Sulfite anions SO_3^{2-}			
1. BaCl_2 (group reagent) in neutral medium gives precipitate BaSO_3 , soluble in strong mineral acids (excluding H_2SO_4). For example, the precipitate dissolves in dilute HCl, when SO_2 (has odour of burning matches) evolves.	Add 1-2 drops of BaCl_2 solution to 2-3 drops of Na_2SO_3 . Divide the sediment on two test tubes. Add 10-15 HCl drops to the first one, and 5-10 NaOH drops to the second one.		$\text{Na}_2\text{SO}_3 + \text{BaCl}_2 = \text{BaSO}_3\downarrow + 2\text{NaCl}$; $\text{SO}_3^{2-} + \text{Ba}^{2+} = \text{BaSO}_3\downarrow$; $\text{BaSO}_3\downarrow + 2\text{HCl} \rightarrow \text{BaCl}_2 + \text{SO}_2\uparrow + \text{H}_2\text{O}$; $\text{BaSO}_3\downarrow + 2\text{H}^+ \rightarrow \text{Ba}^{2+} + \text{SO}_2\uparrow + \text{H}_2\text{O}$. $\text{BaSO}_3\downarrow + \text{NaOH} \rightarrow \text{no react}$

<p>2. Dilute hydrochloric acid HCl or dilute sulfate acid H₂SO₄: decomposition, more rapidly at warming, with the evolution of sulfur oxide(IV) SO₂: The gas may be identified by its suffocating odour of burning sulfur, or by the green colouration, due to the formation of chromium(III) ions, produced when a filter paper, moistened with acidified potassium dichromate solution, is held over the mouth of the test-tube.</p>	<p>Place 8-10 drops of fresh-prepared solution of Sodium Sulfite Na₂SO₃ upon test-tube, add 8-10 drops of HCl. Identify the gas using described method.</p>		$\text{Na}_2\text{SO}_3 + 2\text{HCl} = \text{SO}_2\uparrow + 2\text{NaCl} + \text{H}_2\text{O};$ $\text{SO}_3^{2-} + 2\text{H}^+ = \text{H}_2\text{SO}_3;$ $\text{H}_2\text{SO}_3 \rightarrow \text{SO}_2\uparrow + \text{H}_2\text{O};$ $3\text{SO}_2 + \text{K}_2\text{Cr}_2\text{O}_7 + \text{H}_2\text{SO}_4 = \text{Cr}_2(\text{SO}_4)_3 + \text{K}_2\text{SO}_4 + \text{H}_2\text{O};$ $3\text{SO}_2 + \text{Cr}_2\text{O}_7^{2-} + 2\text{H}^+ = 2\text{Cr}^{3+} + 3\text{SO}_4^{2-} + \text{H}_2\text{O}.$
<p>3. Lime water Ca(OH)₂. This test is carried out by adding dilute hydrochloric acid to the solid sulfite, and bubbling the evolved gas through lime water; a white precipitate of calcium sulfite CaSO₃ is formed.</p>	<p>This test is best carried out in the apparatus shown in fig. The solid Na₂SO₃ (near 0,05-0,1 g) is placed in the test-tube or small distilling flask (10-25 mL capacity), dilute hydrochloric acid added, and the cork immediately replaced. The gas which is evolved (warming may be necessary) is passed into lime water (reagent - in lab fume hood) contained in the test-</p>		$\text{Na}_2\text{SO}_3 + \text{Ca}(\text{OH})_2 = \text{CaSO}_3\downarrow + 2\text{NaOH};$ $\text{SO}_3^{2-} + \text{Ca}^{2+} = \text{CaSO}_3\downarrow$

	<p>tube; the production of a turbidity indicates the presence of a sulfite.</p> 		
<p>4. Strong oxidation agents ($KMnO_4$ or free I_2 solution). Potassium permanganate solution, acidified with dilute sulfate acid before the testing, reacts with Sulfite solution with discolouration, owing to reduction to Mn^{2+} ions. If to use free iodine solution, visual effect of reaction is to discolouration of solution owing to transformation of free I_2 to colourless iodide-ions.</p>	<p>Add 2 drops of Potassium permanganate upon test-tube, acidify by adding of 2-3 drops of H_2SO_4 and add 5-6 drops of fresh-prepared solution of Na_2SO_3. To repeat the previous test with iodine (iodine water) solution.</p>		$2KMnO_4 + 5Na_2SO_3 + 8H_2SO_4 = 2MnSO_4 + 5Na_2SO_4 + K_2SO_4 + 8H_2O;$ $5SO_3^{2-} + 2MnO_4^- + 6H^+ = 2Mn^{2+} + 5SO_4^{2-} + 3H_2O;$ <p>or</p> $2Na_2SO_3 + I_2 = Na_2S_2O_6 + 2NaI;$ $2SO_3^{2-} + I_2 = S_2O_6^{2-} + 2I^- .$

Characteristic reactions of Carbonate anions CO₃²⁻			
1. BaCl₂ (group reagent). Obtained precipitate BaCO ₃ , soluble in HCl with gas bubbles of CO ₂ (no odour) isolation indicates the presence of CO ₃ ²⁻ ions in solution.	Add 2-3 drops of Na ₂ CO ₃ to 2-3 drops of BaCl ₂ . Dilute the obtained sediment in HCl.		Na ₂ CO ₃ + BaCl ₂ = BaCO ₃ ↓ + 2NaCl; CO ₃ ²⁻ + Ba ²⁺ = BaCO ₃ ↓; BaCO ₃ ↓ + 2HCl = BaCl ₂ + CO ₂ ↑ + H ₂ O; BaCO ₃ ↓ + 2H ⁺ = Ba ²⁺ + CO ₂ ↑ + H ₂ O.
2. Mineral acids (H₂SO₄, HCl, HNO₃ etc.), and acetic acid CH₃COOH decompose carbonates with gassing of CO ₂ .	Add 1-2 mL of HCl to 10-15 drops of Na ₂ CO ₃ .		Na ₂ CO ₃ + 2HCl = CO ₂ ↑ + 2NaCl + H ₂ O; CO ₃ ²⁻ + 2H ⁺ = CO ₂ ↑ + H ₂ O;
Characteristic reactions of Phosphate anions PO₄³⁻			
1. BaCl₂ (group reagent) in neutral medium sediments PO ₄ ³⁻ in form of precipitate Ba ₃ (PO ₄) ₂ , soluble in strong acids (except H ₂ SO ₄).	Add 2-3 drops of Na ₂ HPO ₄ to 2-3 drops of BaCl ₂ . Dissolve the obtained precipitate by adding 10-15 drops of HCl.		2Na ₂ HPO ₄ + 3BaCl ₂ = Ba ₃ (PO ₄) ₂ ↓ + 4NaCl + 2HCl; 2HPO ₄ ²⁻ + 3Ba ²⁺ = Ba ₃ (PO ₄) ₂ ↓ + 2H ⁺ ; Ba ₃ (PO ₄) ₂ ↓ + 6HCl = 3BaCl ₂ + 2H ₃ PO ₄ ; Ba ₃ (PO ₄) ₂ ↓ + 6H ⁺ = 3Ba ²⁺ + 2H ₃ PO ₄ .
2. Magnesia mixture (MgCl₂ in presence of buffer mixture NH₃·H₂O + NH₄Cl) reacts with phosphates forming white crystalline precipitate	Add to test tube 2-3 drops of MgCl ₂ and NH ₄ Cl solutions, add to prepared mixture 2-3 drops of (NH ₄) ₂ HPO ₄		Na ₂ HPO ₄ + MgCl ₂ + NH ₃ ·H ₂ O = MgNH ₄ PO ₄ ↓ + 2NaCl + H ₂ O; HPO ₄ ²⁻ + Mg ²⁺ + NH ₃ ·H ₂ O = MgNH ₄ PO ₄ ↓ + H ₂ O.

MgNH ₄ PO ₄ .	solution. Shuffle mixture and add solution of NH ₃ ·H ₂ O for alkali reaction of medium (use litmus).		
3. Molybdenum liquid (the mixture of (NH₄)₂MoO₄ and HNO₃). As the result of reaction the yellow crystalline precipitate (NH ₄) ₃ H ₄ [P(Mo ₂ O ₇) ₆] is formed.	Add 5-8 drops of Molybdenum liquid to 2-3 drops of Phosphate solution. Boil mixture.		Na ₂ HPO ₄ +12(NH ₄) ₂ MoO ₄ +23HNO ₃ = (NH ₄) ₃ H ₄ [P(Mo ₂ O ₇) ₆]↓+ 2NaNO ₃ +21NH ₄ NO ₃ +10H ₂ O; HPO ₄ ²⁻ +3NH ₄ ⁺ +12MoO ₄ ²⁻ +23H ⁺ = =(NH ₄) ₃ H ₄ [P(Mo ₂ O ₇) ₆]↓ +10H ₂ O.
Characteristic reactions of Borate anions B₄O₇²⁻ (BO₂⁻)			
1. BaCl₂ (group reagent) precipitates Barium metaborate Ba(BO ₂) ₂ from concentrated solution of borates.	Add 5-6 drops of BaCl ₂ to 10-15 drops of Na ₂ B ₄ O ₇ (reagent - in lab fume hood). Dissolve the precipitate by adding hydrochloric acid HCl by drops.		Na ₂ B ₄ O ₇ + 2 BaCl ₂ + H ₂ O = 2Ba(BO ₂) ₂ ↓ + 2NaCl + 2HCl; B ₄ O ₇ ²⁻ + 2Ba ²⁺ + H ₂ O = 2Ba(BO ₂) ₂ ↓ + 2H ⁺ ; Ba(BO ₂) ₂ ↓ + 2HCl + 2H ₂ O = BaCl ₂ + 2H ₃ BO ₃ ; Ba(BO ₂) ₂ ↓ + 2H ⁺ + 2H ₂ O = Ba ²⁺ + 2H ₃ BO ₃ .
2. Silver nitrate AgNO₃ precipitates Silver Metaborate from moderate concentrated solutions. In very diluted	Add 1-2 drops of Silver Nitrate solution to 10-15 drops of Na ₂ B ₄ O ₇ (reagent - in lab fume		Na ₂ B ₄ O ₇ + 2AgNO ₃ + 3H ₂ O = 2AgBO ₂ ↓ + 2H ₃ BO ₃ + 2NaNO ₃ . B ₄ O ₇ ²⁻ + 2Ag ⁺ + 3H ₂ O = 2AgBO ₂ ↓ + 2H ₃ BO ₃ .

<p>solutions at cool and in more concentrated ones at heating it forms brown precipitate of Silver Oxide</p> $2\text{AgBO}_2 + 3\text{H}_2\text{O} = \text{Ag}_2\text{O}\downarrow + 2\text{H}_3\text{BO}_3.$	<p>hood). Observe slow formation of precipitate.</p>		
<p>3. Flame test. Bringing of volatile Boron compounds (Methyl Ether $(\text{CH}_3\text{O})_3\text{B}$ or Ethyl Ether $(\text{C}_2\text{H}_5\text{O})_3\text{B}$ or free Borate acid) in colourless flame, it is observed specific colouring of flame.</p>	<p>0,5-1 mL of $\text{Na}_2\text{B}_4\text{O}_7$ (reagent - in lab fume hood) add in small evaporating dish and heat to dry residue on electric stove. Add 10-15 drops of concentrated Sulfate acid to dry residue (Carefully! reagent - in lab fume hood), 1-2 mL of ethyl alcohol (reagent - in lab fume hood) and set fire to surface of mixture in the plate.</p>		$\text{Na}_2\text{B}_4\text{O}_7 + \text{H}_2\text{SO}_4 (\text{conc.}) + 5\text{H}_2\text{O} = 4\text{H}_3\text{BO}_3 + \text{Na}_2\text{SO}_4;$ $\text{B}_4\text{O}_7^{2-} + 2\text{H}^+ + 5\text{H}_2\text{O} = 4\text{H}_3\text{BO}_3;$ $\text{H}_3\text{BO}_3 + 3\text{C}_2\text{H}_5\text{OH} = \text{B}(\text{OC}_2\text{H}_5)_3 + 3\text{H}_2\text{O};$ $2\text{B}(\text{OC}_2\text{H}_5)_3 + 18\text{O}_2 \xrightarrow{t} \text{B}_2\text{O}_3 + 12\text{CO}_2 + 15\text{H}_2\text{O}.$
<p>Characteristic reactions of Molybdate anions MoO_4^{2-}</p>			
<p>1. BaCl_2 (group reagent) precipitates Barium Molybdate BaMoO_4 from concentrated solution of molybdates.</p>	<p>Add 1-2 drops of Silver Nitrate solution to 10-15 drops of Na_2MoO_4 (reagent - in lab fume hood). Weigh a few</p>		$\text{Na}_2\text{MoO}_4 + \text{BaCl}_2 = \text{BaMoO}_4\downarrow + 2\text{NaCl};$ $\text{MoO}_4^{2-} + \text{Ba}^{2+} = \text{BaMoO}_4\downarrow.$

	minutes. Observe slow formation of precipitate.		
2. Ammonium Thiocyanide (Rodanide) NH_4SCN forms complex rodanides with molybdates. Ions Fe^{3+} interferes to this reaction. Fe^{3+} ions are masked by reducing to Fe^{2+} added reducing agents, for example, Sn^{2+} compounds. The last reagents in the same time reduce Mo^{6+} to Mo^{5+} .	Acidify 1 mL of Sodium Molybdate Na_2MoO_4 solution (reagent - in lab fume hood) by 0,5 mL of HCl, add 1-2 mL of saturated solution of Tin(II) chloride $SnCl_2$ (reagent - in lab fume hood) and add 5-6 drops of 10% NH_4SCN (in lab fume hood).		$2Na_2MoO_4 + SnCl_2 + 10NH_4SCN + 12HCl = 2Na_2[MoO(SCN)_5] + SnCl_4 + 10NH_4Cl + 6H_2O;$ $2 MoO_4^{2-} + Sn^{2+} + 10SCN^- + 12H^+ = 2[MoO(SCN)_5]^{2-} + Sn^{4+} + 6H_2O.$
3. Hydrogen Peroxide H_2O_2 in presence of concentrated ammonia $NH_3 \cdot H_2O$ gives Permolybdate red coloured solution. Colour is unstable, may be observed during 10-30 sec after Peroxide was added.	Add 1 mL of concentrated Ammonia (in lab fume hood) to 1-2 mL of Na_2MoO_4 solution (in lab fume hood). Add by drops 10% Hydrogen Peroxide H_2O_2 (in lab fume hood).		$Na_2MoO_4 + 2 NH_3 \cdot H_2O \leftrightarrow (NH_4)_2MoO_4 + 2NaOH;$ $(NH_4)_2MoO_4 + H_2O_2 = (NH_4)_2[MoO_5] + H_2O$ $MoO_4^{2-} + H_2O_2 \leftrightarrow [MoO_5]^{2-} + H_2O$ $Mo^{6+} - 2e \rightarrow Mo^{8+} \quad \quad 1$ $2O^{1-} + 2e \rightarrow 2O^{2-} \quad \quad 1$
Characteristic reactions of Silicate anions SiO_3^{2-}			
1. $BaCl_2$ (group reagent) gives Barium Silicate $BaSiO_3$ precipitate from concentrated	Add 5-6 drops of $BaCl_2$ to 10-15 drops of Na_2SiO_3 (reagent - in lab		$Na_2SiO_3 + BaCl_2 = BaSiO_3 \downarrow + 2NaCl;$ $SiO_3^{2-} + Ba^{2+} = BaSiO_3 \downarrow.$

solution of soluble silicates.	fume hood). Wait a few minutes.		
2. Diluted acids (<i>HCl</i>, <i>H₂SO₄</i>) isolate voluminous gelatinous precipitate of silicate acids from concentrated silicate solutions. At certain conditions (high concentration of silicate, slowly adding of mineral acid) this precipitate coagulates completely and don't pour out from tube. If HCl was added quickly to diluted solution of silicate, silicate acids form colloid solution and don't precipitate for a long time.	Add slowly by drops concentrated HCl (in lab fume hood). to 15-20 drops of Na ₂ SiO ₃ (in lab fume hood).		$\text{Na}_2\text{SiO}_3 + 2\text{HCl} = \text{H}_2\text{SiO}_3\downarrow + 2\text{NaCl};$ $\text{SiO}_3^{2-} + 2\text{H}^+ = \text{H}_2\text{SiO}_3\downarrow.$
3. Silver nitrate <i>AgNO₃</i> isolates yellow precipitate of Silver Silicate Ag ₂ SiO ₃ from concentrated silicate solutions.	Add slowly by drops AgNO ₃ solution (in lab fume hood) to 5-10 drops of concentrated Na ₂ SiO ₃ (in lab fume hood).		$\text{Na}_2\text{SiO}_3 + 2\text{AgNO}_3 = \text{Ag}_2\text{SiO}_3\downarrow + 2\text{NaNO}_3;$ $\text{SiO}_3^{2-} + 2\text{Ag}^+ = \text{Ag}_2\text{SiO}_3\downarrow + 2\text{NO}_3^-.$
4. Ammonium Salts (for example, <i>NH₄Cl</i>) isolate gelatinous precipitate of silicate acids more completely, than mineral acids. Silicate acids	Add 3-6 drops of concentrated NH ₄ Cl (in lab fume hood) solution to 5-10 drops of Na ₂ SiO ₃ solution (in lab fume		$\text{Na}_2\text{SiO}_3 + 2\text{NH}_4\text{Cl} + 2\text{H}_2\text{O} =$ $\text{H}_2\text{SiO}_3\downarrow + 2\text{NH}_3\cdot\text{H}_2\text{O} + 2\text{NaCl};$ $\text{SiO}_3^{2-} + 2\text{NH}_4^+ + 2\text{H}_2\text{O} = \text{H}_2\text{SiO}_3\downarrow + 2\text{NH}_3\cdot\text{H}_2\text{O}.$

precipitate more quickly at heating.	hood), heat mixture in a water bath.		
--------------------------------------	--	--	--

1.7.2. The second group of anions

Content	Description	Observation	Explanation
Characteristic reactions of Chloride anions Cl⁻			
1. Silver Nitrate $AgNO_3$ in presence of HNO_3 reacts with Cl^- forming precipitate $AgCl$, which is dissolved in water ammonia $NH_3 \cdot H_2O$. Reaction is accompanied by the formation of soluble complex compound diamminosilver chloride $[Ag(NH_3)_2]Cl$.	Add 1-2 drops of $AgNO_3$ solution (in lab fume hood) to 2-3 drops of HNO_3 and 1-2 drops of $NaCl$ solution. Add concentrated $NH_3 \cdot H_2O$ (in lab fume hood) to the precipitate up to complete dissolving. Add by drops solution of HNO_3 up to precipitate formation (turbidity).		$NaCl + AgNO_3 = AgCl\downarrow + NaNO_3$; $Cl^- + Ag^+ = AgCl\downarrow$; $AgCl\downarrow + 2NH_3 \cdot H_2O = [Ag(NH_3)_2]Cl + 2H_2O$; $AgCl\downarrow + 2NH_3 \cdot H_2O = [Ag(NH_3)_2]^+ + Cl^- + 2H_2O$; $[Ag(NH_3)_2]Cl + 2HNO_3 = AgCl\downarrow + 2NH_4NO_3$; $[Ag(NH_3)_2]^+ + Cl^- + 2H^+ = AgCl\downarrow + 2NH_4^+$.
Characteristic reactions of Bromide anions Br⁻			
1. Silver Nitrate $AgNO_3$ in presence of HNO_3 gives with Br^- precipitate $AgBr$, which is slightly soluble in $NH_3 \cdot H_2O$ in	Add 1-2 drops of $AgNO_3$ solution (in lab fume hood) to 2-3 drops of HNO_3 and 2-3 drops of		$KBr + AgNO_3 = AgBr\downarrow + KNO_3$; $Br^- + Ag^+ = AgBr\downarrow$.


unlike AgCl.	KBr solution. Add concentrated $\text{NH}_3 \cdot \text{H}_2\text{O}$ (in lab fume hood). Study solubility of sediment in concentrated ammonia.		
2. Chloric water Cl_2 (the solution of Cl_2 in H_2O). Cl_2 reacts with bromides displacing free bromine Br_2 from its salts. The isolated Bromine Br_2 is better dissolved in benzene than in H_2O and benzene layer is colouring yellow.	Add 5-6 drops of benzene C_6H_6 (in lab fume hood) and 4-5 drops of chloric water (in lab fume hood) to 6-8 drops of KBr. Shake the mixture. Wait a few seconds till separating of water and benzene layers.		$2\text{KBr} + \text{Cl}_2 \rightarrow \text{Br}_2 + 2\text{KCl};$ $2\text{Br}^- + \text{Cl}_2 \rightarrow \text{Br}_2 + 2\text{Cl}^-.$
Characteristic reactions of Iodide anions I^-			
1. Silver Nitrate AgNO_3 in presence of HNO_3 reacts with I^- given precipitate AgI, which is slightly soluble in $\text{NH}_3 \cdot \text{H}_2\text{O}$ in unlike AgCl.	Add 1-2 drops of AgNO_3 solution (in lab fume hood) to 2-3 drops of HNO_3 and KI solution. Add concentrated $\text{NH}_3 \cdot \text{H}_2\text{O}$ (in lab fume hood). Study solubility of sediment in concentrated ammonia.		$\text{KI} + \text{AgNO}_3 = \text{AgI}\downarrow + \text{KNO}_3;$ $\text{I}^- + \text{Ag}^+ = \text{AgI}\downarrow.$

<p>2. Chloric water Cl_2 (it is the solution of Cl_2 in H_2O). Cl_2 reacts with iodides displacing free Iodine I_2 from its salts. The isolated I_2 is better dissolved in benzene than in H_2O and benzene layer is colouring pink or crimson. In excess of chloric water colour may vanish due to oxidation of free I_2 to iodate(V) acid HIO_3:</p> $\text{I}_2 + 5\text{Cl}_2 + 6\text{H}_2\text{O} = 2\text{HIO}_3 + 10\text{HCl}$	<p>Add 5-6 drops of benzene C_6H_6 (in lab fume hood) and 4-5 drops of chloric water (in lab fume hood) to 6-8 drops of KI. Shake the mixture. Wait a few seconds till separating of water and benzene layers.</p>		$2\text{KI} + \text{Cl}_2 \rightarrow \text{I}_2 + 2\text{KCl};$ $2\text{I}^- + \text{Cl}_2 \rightarrow \text{I}_2 + 2\text{Cl}^-.$				
<p>3. Copper salts (for example, Copper Sulfate CuSO_4) react with I^- given milk-white precipitate CuI. Mixture has brown colour due to free Iodine I_2.</p>	<p>Mix 5-6 drops of KI and 1-2 drops of CuSO_4. Add 2-4 drops $\text{Na}_2\text{S}_2\text{O}_3$ until discolouration of mixture. Note on the colour of CuI precipitate.</p>		$2\text{CuSO}_4 + 4\text{KI} = 2\text{CuI}\downarrow + \text{I}_2 + 2\text{K}_2\text{SO}_4;$ $2\text{Cu}^{2+} + 4\text{I}^- = 2\text{CuI}\downarrow + \text{I}_2.$ <table style="margin-left: auto; margin-right: auto;"> <tr> <td style="padding-right: 10px;">$\text{Cu}^{2+} + 1\text{e}^- \rightarrow \text{Cu}^{1+}$</td> <td style="border-left: 1px solid black; padding-left: 10px;">2</td> </tr> <tr> <td style="padding-right: 10px;">$2\text{I}^- - 2\text{e}^- \rightarrow \text{I}_2$</td> <td style="border-left: 1px solid black; padding-left: 10px;">1</td> </tr> </table>	$\text{Cu}^{2+} + 1\text{e}^- \rightarrow \text{Cu}^{1+}$	2	$2\text{I}^- - 2\text{e}^- \rightarrow \text{I}_2$	1
$\text{Cu}^{2+} + 1\text{e}^- \rightarrow \text{Cu}^{1+}$	2						
$2\text{I}^- - 2\text{e}^- \rightarrow \text{I}_2$	1						

1.7.3. The third group of anions

Content	Description	Observation	Explanation				
Characteristic reactions of Nitrite anions NO_2^-							
<p>1. Potassium Iodide (KI) is oxidized in acetous medium by nitrite ions up to free I_2. Presence of free Iodine may be</p>	<p>Add 1 drop of fresh-prepared starch, 2 drops of CH_3COOH, 1 drop of KI, and 2-3 drops of</p>		$2\text{KNO}_2 + 2\text{KI} + 4\text{CH}_3\text{COOH} =$ $= \text{I}_2 + 2\text{NO} + 4\text{CH}_3\text{COOK} + 2\text{H}_2\text{O}$ <table style="margin-left: auto; margin-right: auto;"> <tr> <td style="padding-right: 10px;">$\text{N}^{3+} + 1\text{e}^- \rightarrow \text{N}^{2+}$</td> <td style="border-left: 1px solid black; padding-left: 10px;">2</td> </tr> <tr> <td style="padding-right: 10px;">$2\text{I}^- - 2\text{e}^- \rightarrow \text{I}_2$</td> <td style="border-left: 1px solid black; padding-left: 10px;">1</td> </tr> </table>	$\text{N}^{3+} + 1\text{e}^- \rightarrow \text{N}^{2+}$	2	$2\text{I}^- - 2\text{e}^- \rightarrow \text{I}_2$	1
$\text{N}^{3+} + 1\text{e}^- \rightarrow \text{N}^{2+}$	2						
$2\text{I}^- - 2\text{e}^- \rightarrow \text{I}_2$	1						

determined by adding of starch. Starch is very sensitive and specific indicator for Iodine.	NaNO ₂ or KNO ₂).		$2\text{NO}_2^- + 2\text{I}^- + 4\text{CH}_3\text{COOH} = \text{I}_2 + 2\text{NO} + 4\text{CH}_3\text{COO}^- + 2\text{H}_2\text{O}.$
2. Potassium Permanganate (KMnO₄). In acid medium KMnO ₄ oxidizes the nitrite ions up to nitrate ions. The colour of KMnO ₄ disappears and the solution becomes colourless.	Add 1-2 drops of Potassium Permanganate KMnO ₄ , 1-2 drops of H ₂ SO ₄ to 2-3 drops of Sodium or Potassium Nitrite (NaNO ₂ or KNO ₂).		$5\text{KNO}_2 + 2\text{KMnO}_4 + 3\text{H}_2\text{SO}_4 = 5\text{KNO}_3 + 2\text{MnSO}_4 + \text{K}_2\text{SO}_4 + 3\text{H}_2\text{O}$ $\begin{array}{l} \text{N}^{3+} - 2e \rightarrow \text{N}^{5+} \quad \quad 5 \\ \text{Mn}^{7+} + 5e \rightarrow \text{Mn}^{2+} \quad \quad 2 \end{array}$ $5\text{NO}_2^- + 2\text{MnO}_4^- + 6\text{H}^+ = 5\text{NO}_3^- + 2\text{Mn}^{2+} + 3\text{H}_2\text{O}.$
3. Strong acids decompose Nitrites gives gaseous reddish-brown NO ₂ (sometimes named "foxtail").	Add 2-3 drops of Sulfate acid H ₂ SO ₄ to 2-3 drops of Sodium or Potassium Nitrite (NaNO ₂ or KNO ₂). Heat the mixture in a water bath.		$2\text{NaNO}_2 + \text{H}_2\text{SO}_4 = \text{Na}_2\text{SO}_4 + 2\text{HNO}_2;$ $2\text{NO}_2^- + 2\text{H}^+ = 2\text{HNO}_2;$ $2\text{HNO}_2 = \text{NO}_2\uparrow + \text{NO}\uparrow + \text{H}_2\text{O}.$
Characteristic reactions of Nitrate anions NO₃⁻			
1. Iron(II) Sulfate (FeSO₄) (in excess) in presence of concentrated sulfate acid H ₂ SO ₄ gives complex compound [Fe(NO)]SO ₄ . The reaction is called " brown ring ". Note that the presence of	Add 4-5 drops of saturated solution FeSO ₄ (in lab fume hood) to 4-5 drops of NaNO ₃ , shake mixture. The concentrated H₂SO₄ (in lab fume hood) should be CAREFULLY (Be		$2\text{NaN}^{5+}\text{O}_3 + 6\text{Fe}^{2+}\text{SO}_4 + 4\text{H}_2\text{SO}_4 = 3\text{Fe}_2^{3+}(\text{SO}_4)_3 + 2\text{N}^{2+}\text{O} + \text{Na}_2\text{SO}_4 + 4\text{H}_2\text{O}$ $\begin{array}{l} 2\text{Fe}^{2+} - 2e^- \rightarrow 2\text{Fe}^{3+} \quad \quad 3 \\ \text{N}^{5+} + 3e^- \rightarrow \text{N}^{2+} \quad \quad 2 \end{array}$ $\text{NO} + \text{FeSO}_4 = [\text{Fe}(\text{NO})]\text{SO}_4$

<p>nitrite ions will interfere with this test.</p>	<p>attentive! Dangerous) slowly adding by drops at the test-tube wall (in bending state). Do not mix the solution! Slowly lift test-tube in vertical state. A brown ring will form at the junction of the two layers, indicating the presence of the nitrate ion.</p>		
<p>Characteristic reactions of Acetate anions CH_3COO^-</p>			
<p>1. Strong acids. At HCl or H_2SO_4 action to acetates free acetic acid is isolated which can be identified by odour.</p>	<p>Add 3-4 drops of H_2SO_4 to 5-6 drops of Sodium Acetate CH_3COONa, heat the mixture in a water bath.</p>		$2\text{CH}_3\text{COONa} + \text{H}_2\text{SO}_4 = \text{Na}_2\text{SO}_4 + 2\text{CH}_3\text{COOH}\uparrow;$ $\text{CH}_3\text{COO}^- + \text{H}^+ = \text{CH}_3\text{COOH}\uparrow.$
<p>2. Iron(III) Chloride (FeCl_3) with solution of Acetate gives reddish-brown solution of Iron(III) Acetate. When this solution was diluted and heated, the precipitate of basic salt deposited in the result of hydrolysis intensification.</p>	<p>Add 1-2 drops of FeCl_3 solution to 5-6 drops of CH_3COONa. Add 4-6 mL of DW and heat mixture in a water bath until forming of gelatinous precipitate.</p>		$\text{FeCl}_3 + 3\text{CH}_3\text{COONa} = \text{Fe}(\text{CH}_3\text{COO})_3 + 3\text{NaCl};$ $\text{Fe}(\text{CH}_3\text{COO})_3 + 2\text{H}_2\text{O} = \text{Fe}(\text{OH})_2\text{CH}_3\text{COO}\downarrow + 2\text{CH}_3\text{COOH}.$

1.8. PROTOCOL CATIONS MIXTURE OF ANALYSIS

Content of operation	Observation	Conclusion
1. Pre-definition		
A. Colour of solution <i>Previous hypothesis about presence or absence of coloured ions (Fe^{2+}, Fe^{3+}, Cu^{2+}, Co^{2+}, Ni^{2+}, Mn^{2+}) in examined mixture</i>	<i>Your result</i>	_____
B. Odour <i>Previous hypothesis about presence or absence of NH_4^+ ions</i>	<i>Your result</i>	_____
C. pH of solution <i>Previous hypothesis about presence or absence of ions of weak bases (NH_4^+, Fe^{2+}, Fe^{3+}, Cu^{2+}, Co^{2+}, Ni^{2+}, Mn^{2+}, Al^{3+}, Zn^{2+}, Mg^{2+}) in the case of pH less than 7.</i>	<i>Your result</i>	_____
2. Fractional analysis		
1. Test of NH_4^+: To 6-10 drops of the examined mixture add 10 drops of 20% NaOH (in lab fume hood) and the obtained solution heat in a water bath. If ammonia odour fixes or wet litmus paper turns blue, this is NH_4^+ .	<i>Your result</i>	<hr/> <i>Presence or absence NH_4^+ in examined mixture</i>
2. Test of Na^+: To 3-4 drops of the examined mixture add 2 drops of DW and 3-5 drops of Zinc-Uranyl-Acetate $Zn(UO_2)_3(CH_3COO)_8$ (in lab fume hood). Isolation of light-green precipitate means that Na^+ presents in a mixture.	<i>Your result</i>	<hr/> <i>Presence or absence Na^+ in examined mixture</i>

<p>3. Test of Fe^{2+}: To 4 drops of the examined mixture add 2 drops CH_3COOH (pH~5) and 3 drops of $\text{K}_3[\text{Fe}(\text{CN})_6]$. Isolation of dark-blue precipitate means that Fe^{2+} presents in a mixture.</p>	<i>Your result</i>	<hr/> <i>Presence or absence Fe^{2+} in examined mixture</i> <hr/>
<p>4. Test of Fe^{3+}: To 4-5 drops of the examined mixture add 1-2 drops of HCl and 2-3 drops of $\text{K}_4[\text{Fe}(\text{CN})_6]$. Isolation of blue precipitate means that Fe^{3+} presents in a mixture.</p>	<i>Your result</i>	<hr/> <i>Presence or absence Fe^{3+} in examined mixture</i> <hr/>
<p>5. Test of Cu^{2+}: To 12-15 drops of the examined mixture add 10-12 drops of concentrated $\text{NH}_3 \cdot \text{H}_2\text{O}$ (in lab fume hood). Wait 2-3 min. If was observed cornflower color of upper layer of tube content, it means that examined mixture contains Cu^{2+} cations.</p>	<i>Your result</i>	<hr/> <i>Presence or absence Cu^{2+} in examined mixture</i> <hr/>
<p>6. Test of Co^{2+}: Select <i>Variant A</i> or <i>Variant B</i> depends on result, obtained of p. 4.</p> <p>Variant A. In presence of Fe^{3+} (see p. 4): to 6-8 drops of the examined mixture add a few grains of solid NaF; mix content to its complete dissolving. Then add 2-3 drops of NH_4SCN solution, saturated in acetone (in lab fume hood) (if the solution is red, one should add NaF till the red colour disappears). If the blue colour solution is observed - there is Co^{2+}.</p> <p>Variant B. In absence of Fe^{3+} (see p. 4): Add 6-8 drops of the examined mixture to 2-3 drops of NH_4SCN solution saturated in acetone (in lab fume hood). If observes blue colour of solution, then there is Co^{2+}.</p>	<i>Your result</i>	<hr/> <i>Presence or absence Co^{2+} in examined mixture</i> <hr/>

<p>7. Test of Mn^{2+}: To 5-6 drops of the examined mixture add 10 drops of HNO_3 and a few grains of $NaBiO_3$ (in lab fume hood). Wait 1 min. If magenta colour is observed – there is Mn^{2+}.</p>	<i>Your result</i>	<hr/> <i>Presence or absence Mn^{2+} in examined mixture or observe trace quantities of this ion</i>
<p>8. Test of Ni^{2+}: Select <i>Variants A, or B, or C, or D, or E</i> according to results of pp. 3, 4, 5 and 6.</p> <p>Variant A. It is realized in absence Fe^{3+} and Cu^{2+}: To 6-8 drops of the examined mixture add 10-15 drops of concentrated $NH_3 \cdot H_2O$ (in lab fume hood), shuffle tube content and add 2-3 drops of Dimethylglyoxime $C_4H_8N_2O_2$ (Chugaev's reagent) (in lab fume hood); if crimson coloured sediment formed, then there is Ni^{2+}.</p> <p>Variant B. It is realized in presence of Fe^{3+}: To 6-8 drops of the examined mixture add a few grains of NaF up to complete solving, and then 10-12 drops of concentrated $NH_3 \cdot H_2O$ (in lab fume hood) and 2-3 drops of Dimethylglyoxime (in lab fume hood); if crimson precipitate observed, then there is Ni^{2+}.</p> <p>Variant C. It is realized in presence of Cu^{2+}: To 6-8 drops of the examined mixture in centrifuge tube add 3-4 drops of diluted CH_3COOH and then 5-6 drops of KI solution and the precipitate is separated by centrifuge. Add to brown supernatant liquid by drops $Na_2S_2O_3$ till the solution gets colourless. Then one should add 10-15 drops of concentrated $NH_3 \cdot H_2O$ (in lab fume hood) and 2-3 drops of Dimethylglyoxime (in lab fume hood). If crimson coloured sediment observed, then there is Ni^{2+}.</p>	<i>Your result</i>	<hr/> <i>Presence or absence Ni^{2+} in examined mixture or observe trace quantities of this ion</i>

<p>Variant D. It is realized in presence of Fe³⁺ and Cu²⁺ at the same time. To 6-8 drops of the examined mixture in centrifuge tube add 3-4 drops of diluted CH₃COOH and then 5-6 drops of KI solution and the precipitate is separated by centrifuge. Add to brown supernatant liquid by 5-10 drops of Na₂S₂O₃. Then a few grains of NaF up to complete solving, and then 10-12 drops of concentrated NH₃·H₂O (in lab fume hood) and 2-3 drops of Dimethylglyoxime (in lab fume hood); if crimson precipitate observed, then there is Ni²⁺.</p> <p>Variant E. It is realized in presence of Fe²⁺, it is oxidized by 2-3 drops of the 6% solution of H₂O₂ (in lab fume hood) to Fe³⁺, and then add a few grains of NaF up to complete solving, and then 10-12 drops of concentrated NH₃·H₂O (in lab fume hood) and 2-3 drops of Dimethylglyoxime (in lab fume hood); if crimson coloured sediment, then there is Ni²⁺.</p>		
3. Systematic analysis		
<p>9. Test for K⁺: Select Variant A or Variant B according to results of pp. 1, 5.</p> <p>Variant A realizes in absence of NH₄⁺ and Cu²⁺: To 1-2 mL of examined mixture (in centrifugal test-tube) add 1-2 mL of saturated solution of Na₂CO₃ (in lab fume hood); the mixture is centrifugated, supernatant liquid is tested for fullness of sediment. Then one should add 1-2 drops of CH₃COOH to pH = 6, to the obtained solution one adds 5-6 drops of Na₃[Co(NO₂)₆] (in lab fume hood); if there is yellow-orange precipitate - then there is K⁺.</p>		

Variant B. It is realized **in presence of NH_4^+ , but absence of Cu^{2+}** : To 2-3 mL of the examined mixture (in centrifugal test-tube) add by drops 2-3 mL of saturated solution of Na_2CO_3 (**in lab fume hood**); the mixture is centrifugated. The filtrate is transported in the crucible, evaporate liquid, and then calcinate in muffle during 1,5 hours at 700-800 °C. A few crystals of dry residue transport in test-tube, add 10 drops of DW and 1-2 drops of Nessler's reactant (**in lab fume hood**). If no reddish precipitate, NH_4^+ was removed completely. Residue is dissolved in 1-2 mL of DW. Then add 1-2 drops CH_3COOH to pH=6, 5-6 drops of $\text{Na}_3[\text{Co}(\text{NO}_2)_6]$ (**in lab fume hood**); if there is yellow-orange precipitate - there is K^+ .

Variant C. It is realized **in absence of NH_4^+ , but presence of Cu^{2+}** : To 1-2 mL of the examined mixture in centrifugal test-tube add 7-10 drops of solution of NaI (**in lab fume hood**), and the precipitate obtained is separated using centrifuge (don't forget test of fullness of precipitation!). The supernatant solution has to be decanted in test-tube. Then $\text{Na}_2\text{S}_2\text{O}_3$ is added till the solution gets colourless. Add 1-2 mL of saturated solution of Na_2CO_3 (**in lab fume hood**). The mixture is centrifugated; supernatant liquid is tested for fullness of sedimentation. Then one should add 1-2 drops of CH_3COOH to pH = 6, to the obtained solution one adds 5-6 drops of $\text{Na}_3[\text{Co}(\text{NO}_2)_6]$ (**in lab fume hood**); if there is yellow-orange precipitate - then there is K^+ .

Your result

Presence or absence K^+ in examined mixture

<p>10. Separation and tests of the 4th group cations (Ag^+ and Pb^{2+}): To 2 drops of the examined mixture add 2 drops of HCl.</p> <p>- If no precipitate, there is no Ag^+ and/or Pb^{2+} in examined mixture. Go to p. 11.</p> <p>If precipitate was observed, the order of operation is the next. To 2-3 mL of the examined mixture in centrifugal tube add 2-3 mL of 10% HCl (in lab fume hood); the sediment obtained is separated using centrifuge (don't forget test of fullness of precipitation). The upper liquid is decanted in separate labeled tube and stores it for the next analysis (Filtrate 1). Solid residue – Precipitate 1 in centrifuge test-tube is rinsed with 2-3 mL of hot DW and is centrifuged again (for separating of PbCl_2 soluble in hot water), which gives us the Filtrate 2 and the Precipitate 2. The Filtrate 2 is decanted into the clean test-tube; then add 3-4 drops KI. If forms yellow precipitate, then there is Pb^{2+}.</p> <p>If the Precipitate 2 didn't dissolved completely in hot water, it should be processed by adding 1-2 mL of concentrated $\text{NH}_3 \cdot \text{H}_2\text{O}$ solution (in lab fume hood), and centrifuged again. The Filtrate 3 is decanted in a clean test-tube; then add HNO_3 solution to $\text{pH} = 3-4$ (using the universal litmus indicator). If the turbidity or bulky white precipitate observes, then there is Ag^+.</p>	<p><i>Your result</i></p>	<hr/> <p><i>Presence or absence Ag^+ and/or Pb^{2+} in examined mixture</i></p>
---	---------------------------	---

<p>11. Separation and tests of cations Ca^{2+} and Sr^{2+}: Select Variant A or B according to result of p. 10.</p> <p>Variant A – if no detected the presence of the 4th group cations (Ag^+ and/or Pb^{2+}). To 5-10 drops of examined mixture add 5-6 drops of 20% H_2SO_4 (in lab fume hood) and 2-3 drops of acetone (or alcohol) (in lab fume hood). If there is precipitate, then there is Ca^{2+} and/or Sr^{2+}. If the solution remains clean, then there is no Ca^{2+} and Sr^{2+}. Go to p. 12.</p> <p>Test of Ca^{2+}. In centrifuge test tube add 1-2 mL of 20% H_2SO_4 (in lab fume hood) and 5-10 drops of acetone (in lab fume hood) to 2-3 mL of the examined mixture. Content of test-tube should be mixed and heated in a water bath for 3-4 min. (to 70-80°C). The mixture is cooled down, and centrifuged, obtained the Precipitate 4 and the Filtrate 4 (store it in the separate labeled tube for the next analysis). To the Precipitate 4 in centrifuge test-tube add 2-3 mL of concentrated $(\text{NH}_4)_2\text{SO}_4$ solution (in lab fume hood) and then heat it in the water bath 3-5 min. If the precipitate didn't dissolve completely, then it should be centrifuged again. Decant the Filtrate 5 in clean test-tube, add by drops $\text{NH}_3 \cdot \text{H}_2\text{O}$ to pH ~ 5-6 (by universal litmus indicator) and add 5-6 drops of $(\text{NH}_4)_2\text{C}_2\text{O}_4$. If there is white precipitate, then there is Ca^{2+}.</p> <p>Test of Sr^{2+}: Add in test-tube 1-2 mL of saturated $(\text{NH}_4)_2\text{SO}_4$ solution (in lab fume hood) and heat 1-2 min. in a water bath, and then add 10-12 drops of the examined mixture. If there is white precipitate, it means that Sr^{2+} presents in the mixture.</p>	<p><i>Your result</i></p>	<hr/> <p><i>Present or absent Ca^{2+} and/or Sr^{2+} in examined mixture</i></p>
--	---------------------------	--

<p>Variante B. It is realized in presence of the 4th group cations (Ag^+ and/or Pb^{2+}) like the Variante A, there is one difference only – examined mixture is replaced of the Filtrate 1 (was obtained in p. 11 and was stored in labeled tube) .</p>		
<p>12. Separation and tests of cations Al^{3+} and Zn^{2+}: Select Variante A, B, or C depends on results of p. 7, 10 and 11.</p> <p>Variante A. If detected the absence of Mn^{2+} (see p. 7), absence the 4th group cations (Ag^+ and Pb^{2+}) (see p. 10), absence cations Ca^{2+} and Sr^{2+} (see p. 11). In centrifuge test tube mix 2-3 mL of examined mixture and 1-2 mL of 10% NaOH (in lab fume hood), heat in a water bath for 2-3 min. After cooling this mixture it should be centrifuged, and obtained The Precipitate 6 (don't use in the next operations) and The Filtrate 6. Decant Filtrate 6 in two test-tubes.</p> <p>Test of Al^{3+}: To the Filtrate 6 in the first test tube add a few crystals of solid NH_4Cl till getting the saturated solution. Heat a test tube in a water bath (for 2-3 min.). If white gelatinous precipitate $\text{Al}(\text{OH})_3$ was observed, it means that Al^{3+} was contained in mixture.</p> <p>Test of Zn^{2+}: To the Filtrate 6 in the second test tube add by drops HCl (to pH~5 by universal litmus paper) and 5-6 drops of $\text{K}_3[\text{Fe}(\text{CN})_6]$. If there is yellow-and-orange precipitate, then there is Zn^{2+}.</p>	<p><i>Your result</i></p>	<hr/> <p><i>Presence or absence Al^{3+} and/or Zn^{2+} in examined mixture</i></p>

Variant B. If detected the presence of Mn^{2+} (see p. 7), but absence the 4th group cations (Ag^+ and Pb^{2+}) (see p. 10), and absence cations Ca^{2+} and Sr^{2+} (see p. 11). In centrifuge test tube mix 2-3 mL of examined mixture, 5-6 drops of 6% H_2O_2 (in lab fume hood), and 1-2 mL of 10% NaOH (in lab fume hood), heat in a water bath for 2-3 min. Cool this mixture, centrifuge, and obtained **The Precipitate 6** (don't use in the next operations) and **The Filtrate 6**. Decant Filtrate 6 in two test-tubes. Continue as Variant A – tests of Al^{3+} and Zn^{2+} .

Variant C. It is realized in the presence of Mn^{2+} (see p. 7), presence of the 4th group (Ag^+ and/or Pb^{2+}) (see p. 10), but absence of Ca^{2+} and/or Sr^{2+} . In centrifuge test tube mix 1-2 mL of the **Filtrate 1**, 5-6 drops of 6% H_2O_2 (in lab fume hood), and 1-2 mL of 10% NaOH (in lab fume hood), heat in a water bath for 2-3 min. After cooling this mixture it should be centrifuged, and obtained **The Precipitate 6** (keep away) and **The Filtrate 6**. Decant Filtrate 6 in two test-tubes. Continue as Variant A – tests of Al^{3+} and Zn^{2+} .

Variant D. It is realized in the presence of Mn^{2+} (see p. 7), presence of the 4th group (Ag^+ and/or Pb^{2+}) (see p. 10), and presence of Ca^{2+} and/or Sr^{2+} . In centrifuge test tube mix 1-2 mL of the **Filtrate 4**, 5-6 drops of 6% H_2O_2 (in lab fume hood), and 1-2 mL of 10% NaOH (in lab fume hood), heat in a water bath for 2-3 min. After cooling this mixture it should be centrifuged, and obtained **The Precipitate 6** (keep away) and **The Filtrate 6**. Decant Filtrate 6 in two test-tubes. Continue as Variant A – tests of Al^{3+} and Zn^{2+} .

Variant E. It is realized **in the absence of Mn^{2+}** (see p. 7), **presence of the 4th group (Ag^+ and/or Pb^{2+})** (see p. 10), and **absence of Ca^{2+} and/or Sr^{2+}** . In centrifuge test tube mix 1-2 mL of **the Filtrate 1** and 1-2 mL of 10% NaOH (**in lab fume hood**), heat in a water bath for 2-3 min. After cooling this mixture it should be centrifuged, and obtained **The Precipitate 6** (don't use in the next operations) and **The Filtrate 6**. Decant Filtrate 6 in two test-tubes. Continue as Variant A – tests of Al^{3+} and Zn^{2+} .

Variant F. It is realized **in the absence of Mn^{2+}** (see p. 7), **absence of the 4th group (Ag^+ and/or Pb^{2+})** (see p. 10), **but presence of Ca^{2+} and/or Sr^{2+}** . In centrifuge test tube mix 1-2 mL of **the Filtrate 4** and 1-2 mL of 10% NaOH (**in lab fume hood**), heat in a water bath for 2-3 min. After cooling this mixture it should be centrifuged, and obtained **The Precipitate 6** (don't use in the next operations) and **The Filtrate 6**. Decant Filtrate 6 in two test-tubes. Continue as Variant A – tests of Al^{3+} and Zn^{2+} .

Variant G. It is realized **in the presence of Mn^{2+}** (see p. 7), **absence of the 4th group (Ag^+ and/or Pb^{2+})** (see p. 10), and **presence of Ca^{2+} and/or Sr^{2+}** . In centrifuge test tube mix 1-2 mL of **the Filtrate 4**, 5-6 drops of 6% H_2O_2 (**in lab fume hood**), and 1-2 mL of 10% NaOH (**in lab fume hood**), heat in a water bath for 2-3 min. After cooling this mixture it should be centrifuged, and obtained **The Precipitate 6** (don't use in the next operations) and **The Filtrate 6**. Decant Filtrate 6 in two test-tubes. Continue as Variant A – tests of Al^{3+} and Zn^{2+} .

<p>Variante G. It is realized in the absence of Mn^{2+} (see p. 7), presence of the 4th group (Ag^+ and/or Pb^{2+}) (see p. 10), and presence of Ca^{2+} and/or Sr^{2+}. In centrifuge test tube mix 1-2 mL of the Filtrate 4, and 1-2 mL of 10% NaOH (in lab fume hood), heat in a water bath for 2-3 min. After cooling this mixture it should be centrifuged, and obtained The Precipitate 6 (don't use in the next operations) and The Filtrate 6. Decant Filtrate 6 in two test-tubes. Continue as Variant A – tests of Al^{3+} and Zn^{2+}.</p>		
<p>13. Separation and analysis of cation Mg^{2+}: Select Variant A or B depends on the result of p. 5.</p> <p>Variante A: It is realized in the presence of Cu^{2+} (see p. 5) in the examined mixture. In centrifuge test-tube add 8-10 drops of NaI (in lab fume hood) to 1-2 mL of the examined mixture The sediment obtained should be centrifuged: there is the precipitate (don't use in the next operations) and The filtrate 7.</p> <p>Decant the Filtrate 7 in a centrifuge test-tube, add by drops of $Na_2S_2O_3$ till the solution gets colourless, then one should of the 5-6 drops of solution NH_4Cl, 5-6 drops of concentrated $NH_3 \cdot H_2O$ (in lab fume hood) and 15-20 drops of concentrated $(NH_4)_2CO_3$ (in lab fume hood). The mixture is heated in a water bath 2-3 min, cooled after that, and centrifuged. The precipitate (don't use in the next operations) and the Filtrate 8 are obtained. To the Filtrate 8 is decanted in a test-tube, add 5-6 drops of Na_2HPO_4. Formation of white precipitate indicates the presence of Mg^{2+} ions.</p>	<p><i>Your result</i></p>	<hr/> <p><i>Presence or absence Mg^{2+} in examined mixture</i></p>

<p>Variant B: It is realized in the absence of Cu^{2+} (see p. 5): to 1-2 mL of the examined mixture add 5-6 drops of concentrated $\text{NH}_3 \cdot \text{H}_2\text{O}$ (in lab fume hood) and 15-20 drops of concentrated $(\text{NH}_4)_2\text{CO}_3$ (in lab fume hood). The mixture is heated in a water bath 2-3 min, cooled after that, and centrifuged. The precipitate (don't use in the next operations) and the Filtrate 7 are obtained. To the Filtrate 7 is decanted in a test-tube, add 5-6 drops of Na_2HPO_4. Formation of white precipitate indicates the presence of Mg²⁺ ions.</p>		
---	--	--

CONCLUSION: Examined mixture contains cations _____.
(list of detected cations)

1.9. QUALITATIVE ANALYSIS OF THE INORGANIC SUBSTANCES

1.9.1 A soluble salt

Unknown substance is considered as the salt soluble in water, and it consists of one cation and one anion. Tests of cations and anions are executed till the first positive reaction.

Qualitative analysis of unknown substance consists of series of operations:

- A. Pre-definition;
- B. Examination of cation component;
- C. Examination of anion component.

A. Pre-definition

Describe *colour*:

Colourless – research object probably doesn't include coloured cations (Fe^{2+} , Mn^{2+} , Fe^{3+} , Cu^{2+} , Co^{2+} , Ni^{2+});

Fe^{2+} – light-green, light-grey;

Mn^{2+} – slightly pink;

Cu^{2+} – dark blue, blue, green-and-blue;

Co^{2+} – pink, violet;

Ni^{2+} – light green;

Fe^{3+} – yellow-and-brown;

and *relative size* of particles (coarse-crystalline or powder).

Draw attention to *odour* of substance. Ammonium- and acetate-contained salts have a characteristic odour.

Dissolving of unknown salt

The tests for dissolving of the substance are started from the processing of some quantity of its grains by salt, which are taken for the analysis, are easily dissolved in DW or in diluted nitrate acid HNO_3 .

The first testing object, proposed for analysis, is soluble in water or is the solution (in the case of high hygroscopic).

a) If the received sample is the solid substance, so 0,15-0,2 g of substance is dissolved in 6-8 mL of DW (if necessary the solution is heated in a water bath);

b) If the received sample is the solution, so it should be diluted: 3-4 mL of DW is added to 1 mL of solution.

A *pH* of received solution is examined using universal litmus paper. Acidic reaction of solution indicates that the sample can be the salt of weak base and strong acid (for example, AlCl_3 , NH_4Cl , $\text{Zn}(\text{NO}_3)_2$). Alkaline reaction of the solution gives the salts formed of weak acid and strong base

(for example, Na_2CO_3 , K_3PO_4 , CH_3COONa). The neutral reaction of the solution indicates that the examined substance is the salt of strong acid and strong base or weak acid and weak base (for example, KCl , Na_2SO_4 , CaCl_2 , $\text{CH}_3\text{COONH}_4$).

B. Determination of cations

1. Test for the presence of the 4th cations group

Add the group reagent HCl to separate portion of the examined solution. If precipitate was observed so the cation belongs to the 4th group – Ag^+ or Pb^{2+} .

Test of Ag^+ – to precipitate, obtained at HCl acting, add concentrated $\text{NH}_3 \cdot \text{H}_2\text{O}$ (*in lab fume hood*). If precipitate dissolved completely, add nitrate acid HNO_3 by drops till turbidity;

Test of Pb^{2+} – with KI and subsequent re-crystallization of “gold rain”.

2. Test for the presence of the 3^d and 2^d cations groups

Add the group reagent $(\text{NH}_4)_2\text{HPO}_4$ to separate portion of the solution. If precipitate was formed so the cation can belong to 3^d or 2^d group. In this case add some drops of concentrated $\text{NH}_3 \cdot \text{H}_2\text{O}$ (*in lab fume hood*) to the isolated precipitate. If we see that:

- The precipitate dissolved, so the cation belongs to the 3^d group;
- The precipitate is not dissolved so the cation belongs to the 2^d group.

Variant A. The precipitate is dissolved - cation of the 3^d group; the reactions are performed in separate portions of the solution:

Test of Cu^{2+} – with $\text{NH}_3 \cdot \text{H}_2\text{O}$ (excess) (*in lab fume hood*);

Test of Zn^{2+} – with $\text{K}_3[\text{Fe}(\text{CN})_6]$;

Test of Co^{2+} – with NH_4SCN in acetone (*in lab fume hood*);

Test of Ni^{2+} – with of Dimethylglyoxime (Chugaev's reagent) $\text{C}_4\text{H}_8\text{N}_2\text{O}_2$ (*in lab fume hood*).

Variant B. If the precipitate is not dissolved – the 2^d cations group:

Test of Fe^{2+} – with $\text{K}_3[\text{Fe}(\text{CN})_6]$;

Test of Fe^{3+} – with $\text{K}_4[\text{Fe}(\text{CN})_6]$ or NH_4SCN ;

Test of Mn^{2+} – with NaBiO_3 (*in lab fume hood*) + HNO_3 ;

Test of Al^{3+} – with NaOH and crystalline NH_4Cl at heating;

Test of Ca^{2+} – with $(\text{NH}_4)_2\text{C}_2\text{O}_4$;

Test of Sr^{2+} – with gypsum water (saturated water solution of CaSO_4) (*in lab fume hood*);

Test of Mg^{2+} – with Na_2HPO_4 in presence of $\text{NH}_3 \cdot \text{H}_2\text{O}$ + NH_4Cl .

3. Determination of the 1st cations group

The sequence and conditions of reaction performance should be followed:

Test of NH_4^+ – with NaOH at heating;

Test of K^+ – with $\text{Na}_3[\text{Co}(\text{NO}_2)_6]$ (*in lab fume hood*); or $\text{NaHC}_4\text{H}_4\text{O}_6$;

Test of Na^+ – with Zinc-Uranyl-Acetate $\text{Zn}(\text{UO}_2)_3(\text{CH}_3\text{COO})_8$ (*in lab fume hood*).

C. Determination of anions


It is concluded that series of anions can be present in solution after the definition of cation, taking into account that unknown substance is soluble in water.

1. Determination of the 1st anions group

Add BaCl_2 group reagent to the separate portion of examined solution. If the precipitate is formed so the anion is of the first group. Remember, that anions $\text{B}_4\text{O}_7^{2-}$, SiO_3^{2-} , MoO_4^{2-} form precipitates only from concentrated solutions.

Test of SO_4^{2-} – HCl acts to precipitation. If the precipitate is not dissolved so SO_4^{2-} is present in examined substance.

Test of CO_3^{2-} – as the result of HCl action to precipitate, the last is dissolved with bubbles gassing. To be sure of CO_3^{2-} to pass gas through acidified weak solution of KMnO_4 used test system



If permanganate solution keeps light-crimson, CO_3^{2-} is present.

Test of SO_3^{2-} – as the result of HCl action to precipitation, the last is dissolved with bubbles gassing. To be sure of SO_3^{2-} to pass gas through acidified weak solution of KMnO_4 used gas test system (see above). If permanganate solution decolourized, SO_3^{2-} is present.

Test of PO_4^{3-} – as the result of HCl action, the precipitate is dissolved without gas isolation. It can be supposed that there is PO_4^{3-} . Then perform the reaction of PO_4^{3-} determination with molybdenum liquid (*in lab fume hood*) at heating in the separate portion of examined solution.

Even if the result of BaCl_2 action the precipitate is not observed, examine the presence of $\text{B}_4\text{O}_7^{2-}$, SiO_3^{2-} , MoO_4^{2-} .

Test of $\text{B}_4\text{O}_7^{2-}$ – flame test with H_2SO_4 (conc.) (**CAREFULLY!**) (*in lab fume hood*) and alcohol (*in lab fume hood*);

Test of SiO_3^{2-} – with concentrated NH_4Cl solution (*in lab fume hood*);

Test of MoO_4^{2-} – with 10% H_2O_2 (*in lab fume hood*) in presence of concentrated ammonia $\text{NH}_3 \cdot \text{H}_2\text{O}$ (*in lab fume hood*).

If all tests are negative, so there is *no the first group* of anions and the search should be continued in the second group of anions.

2. Determination of the 2^d anions group

Add AgNO_3 group reagent (*in lab fume hood*) to the separate portion of examined solution in presence of HNO_3 . If the precipitate of white (yellow-white) colour is formed so anion of the 2^d group is present.

Test of Cl^- – if the precipitate, obtained on the action of group reagent is dissolved in $\text{NH}_3 \cdot \text{H}_2\text{O}$ and as the result of HNO_3 adding the precipitate becomes turbid so there is Cl^- ion.

Test of Br^- and I^- – chloric water Cl_2 (*in lab fume hood*) and benzene C_6H_6 (*in lab fume hood*) add to the examined solution. If benzene layer becomes yellow, Br^- is present, crimson – I^- is present.

If no sediment as the result of group reagent action – *no the second group* of anions, the search should be continued in the third anion group.

3. Determination of the 3^d anions group

Perform the determination reactions in separate portions of examined solution:

Test of NO_3^- – with saturated FeSO_4 (*in lab fume hood*) + concentrated H_2SO_4 (*in lab fume hood*);

Test of NO_2^- – with $\text{KI} + \text{CH}_3\text{COOH}$;

Test of CH_3COO^- – with HCl at heating.

On the basis of obtained results the report on unknown substance analysis should be prepared and the formula of salt should be given (below).

**PROTOCOL OF ANALYSIS
SUBSTANCE # 1 (Soluble salt)**

Stage of analysis	Content of operation	Observation	Conclusion
1. Predefinition	Colour of solid salt, hygroscopicity		
	Odour		
	Solubility in water		
	pH of water solution		
2. Determination of cations	Test for the presence of the 4th cations group: + HCl <i>Test of Ag⁺</i> _____ <i>Test of Pb²⁺</i> _____		
	Test for the presence of the 3^d and 2^d cations groups (if necessary): + (NH₄)₂HPO₄+ NH₄Cl+NH₃·H₂O; + NH₃·H₂O (in excess) <i>Test of Cu²⁺</i> _____ <i>Test of Zn²⁺</i> _____ <i>Test of Co²⁺</i> _____ <i>Test of Ni²⁺</i> _____		

	<p><i>Text of Fe²⁺</i> _____</p> <p><i>Test of Fe³⁺</i> _____</p> <p><i>Test of Mn²⁺</i> _____</p> <p><i>Test of Al³⁺</i> _____</p> <p><i>Test of Ca²⁺</i> _____</p> <p><i>Test of Sr²⁺</i> _____</p> <p><i>Test of Mg²⁺</i> _____</p>		
	<p>Test of the presence of the 1st cations group (if necessary):</p> <p><i>Test of NH₄⁺</i> _____</p> <p><i>Test of K⁺</i> _____</p> <p><i>Test of Na⁺</i> _____</p>		
3. Determination of anions	<p>Test of the 1st anions group: + BaCl₂</p> <p><i>Test of SO₄²⁻</i> _____</p> <p><i>Test of CO₃²⁻</i> _____</p> <p><i>Test of PO₄³⁻</i> _____</p> <p><i>Test of B₄O₇²⁻</i> _____</p>		

	<i>Test of SiO_3^{2-} _____</i> <i>Test of MoO_4^{2-} _____</i>		
	Test of the 2^d anions group (if necessary): <i>Test of Cl^- _____</i> <i>Test of Br^- or I^- _____</i>		
	Test of the 3^d group of anions (if necessary): <i>Test of NO_3^- _____</i> <i>Test of NO_2^- _____</i> <i>Test of CH_3COO^- _____</i>		
CONCLUSION: Formula of salt is _____			

1.9.2 Insoluble substances

Unknown substance may be insoluble in water.

In the first case, the previous step of analysis – to dissolve of unknown substance.

It may be free metals, oxides, basic salts, carbonates, phosphates. Given such substances will be considered to consist of one cation and one anion. Tests of cations and anions are executed till the first positive reaction. The most of such compounds are dissolving in acids or alkalis.

Test of solubility includes examining of the next solvents:

- Boiled DW (for example, PbCl_2 is more soluble in hot water, than in cool water);
- Diluted acetic acid CH_3COOH – dissolves the carbonates (**with gassing of CO_2**), oxides, hydroxides;
- Diluted nitrate acid HNO_3 – for dissolving of free heavy metals, oxides, basic salts, phosphates, and hydroxides;
- Concentrated nitrate acid HNO_3 – for dissolving of free heavy metals, oxides and hydroxides;
- “Aqua regia” – mixture of concentrated HCl and concentrated HNO_3 (in volume ratio 3:1) – for dissolving of the most insoluble compounds.

The next order of analysis is the same described above. **Remember, that prepared solution would be acidic, if we dissolved the unknown substance in weak or strong acid.** For tests of cations and anions it is necessary to regulate pH of examined solution to add corresponding agents (for example, diluted NaOH by drops till required medium). Of course, anion of acid, used for dissolving, will present in examined solution.

For tests of cations and anions it is necessary to regulate pH of examined solution to add corresponding agents. Remember, that components of analytes used for regulation of pH will be detected in examined solution.

The next order of analysis is the same one as described in p. 1.9.1.

PROTOCOL OF ANALYSIS
SUBSTANCE # 2 (Insoluble)


Stage of analysis	Content of operation	Observation	Conclusion
1. Predefinition	Colour of Substance		
	Solubility in: <ul style="list-style-type: none"> ■ Boiled DW; ■ Diluted acetic acid CH_3COOH; ■ Diluted nitrate acid HNO_3; ■ Concentrated nitrate acid HNO_3; ■ "Aqua regia" Observation of gassing		
	pH of obtained solution		
2. Determination of cations	Test for the presence of the 4 ^d cations group: <i>Test of Ag^+</i> _____ _____ Test for the presence of the 3 ^d and 2 ^d cations groups (if necessary): <i>Test of Cu^{2+}</i> _____ _____ _____ _____		

	<hr/> <hr/> <hr/> <hr/>		
3. Determination of anions	Test of the 1st anions group: <i>Test of SO_4^{2-}</i> <hr/> <hr/> <hr/>		
	Test of the 2^d anions group (if necessary): <i>Test of Cl^-</i> _____ <hr/>		
	Test of the 3^d anions group (if necessary): <i>Test of NO_3^-</i> _____ <hr/> <hr/>		
CONCLUSION: Formula of substance is _____			

EXAMPLES OF CONTROL TASKS FOR QUALITATIVE ANALYSIS

1. Write the equation of the chemical reaction of group reagent $(\text{NH}_4)_2\text{HPO}_4$ (in the presence of ammonia $\text{NH}_3 \cdot \text{H}_2\text{O}$) and soluble Strontium soluble salt (for example, $\text{Sr}(\text{NO}_3)_2$) in molecular form. Write the equation of the dissolving of obtained sediment in the Acetic acid CH_3COOH .

2. Propose the reagent for the separation of cations:



Solution: ___ Cation

Sediment: ___ (chemical formula)

Write the molecular equation of the precipitation reaction.

3. Note on the characteristic analytical test of Molybdate MoO_4^{2-} anions:

A	Action of Hydrogen peroxide H_2O_2 and water ammonia $\text{NH}_3 \cdot \text{H}_2\text{O}$	C	Action of Silver nitrate AgNO_3 and the following dissolving of isolated sediment in water ammonia $\text{NH}_3 \cdot \text{H}_2\text{O}$
B	Action of FeSO_4 (saturated) and H_2SO_4 (concentrated)	D	Action of Potassium Permanganate KMnO_4 in presence of H_2SO_4 (diluted)

4. Write reaction of FeCl_3 and yellow blood salt in molecular form and calculate the sum of coefficients in this reaction (*remember, that absence of coefficient before formula means figure 1*)

5. Note the correspondence of the anions and their group reagents:

A	BaCl_2 (at pH=7)	1	NO_3^-
B	AgNO_3 (at pH=2)	2	CH_3COO^-
C	No group reagent	3	I^-
		4	CO_3^{2-}
		5	Cl^-
		6	PO_4^{3-}

6. Note on the highest pure analyte grade of a reagent:

A	Extra Pure grade	C	Guaranteed Reagent (GR)
B	Ultra pure grade	D	Technical grade

7. Minimum value of concentration limit of qualitative analytical reaction must be not less than

A	1:1000
B	1:100000
C	1:10
D	1:1000000

8. NaOH may be the selective analytical reagent for such group of cations:

A	Ions of alkali metals;
B	Ions of alkali-earth metals;
C	Ions of amphoteric metals;
D	Ions of d-elements.

9. What reactant may be used for separation of Ca^{2+} and Co^{2+} cations?

A	NaOH;
B	$(\text{NH}_4)_2\text{CO}_3$ (concentrated);
C	NH_4Cl (concentrated);
D	$\text{NH}_3 \cdot \text{H}_2\text{O}$ (in excess).

10. Put in the sentence a missing word: *Qualitative reaction of SiO_3^{2-} anion with NH_4Cl (saturated solution) is accompanied by visual effect of _____ sediment.*

11. Determine correspondence of the cations and their groups according to phosphate classification:

A	1st	1	Ca^{2+}
B	2d (A)	2	Cu^{2+}
C	2d (B)	3	NH_4^+
D	3d	4	Fe^{2+}
E	4th	5	Fe^{3+}
		6	Co^{2+}
		7	Mg^{2+}
		8	K^+
		9	Ag^+

CHAPTER 2. QUNTITATIVE ANALYSIS

2.1. Lab work - Precipitate gravimetry (mass analysis). Determination of Barium content in Hydrated Barium Chloride

REAGENTS

1. Sulfate acid H_2SO_4 , 2N solution;
2. Hydrochloric acid HCl , 2N solution;
3. DW.

GLASSWARE

1. Watch glass, 80-90 mm diameter;
2. Two beakers, volume 230-300 mL;
3. Wash bottle;
4. Glass stick;
5. Glass funnel;
6. Porcelain crucible, 30-35 mm diameter;
7. Dessicator;
8. Filter paper (Whatman Blue Ribbon filter paper. Ashless standard filter paper for very fine precipitates (class 2d acc. to DIN 53 135);
9. Scissors;
10. Spatula;
11. Crucible tongs.

EQUIPMENT

1. Analytical balances;
2. Electric oven;
3. Lab dry box;
4. Muffle.

BACKGROUND

The historical method of choice for sulfate in waters and wastewaters is the gravimetry with barium:



1. Barium sulfate precipitates quantitatively by adding excess of ion Ba^{2+} under acidic conditions; low pH is required to avoid the co-precipitation of barium carbonates and/or phosphates.

2. The formation of high purity and non-filterable crystals implies to allow reaction to continue for at least 2 hours at temperature over 80-90 °C. This process (digestion of the precipitate) minimizes the formation of filterable BaSO_4 crystals from the initial colloidal particles.

3. Filter the precipitate by decantation on a filter paper and the resulting solid mass is washed and dried in an oven for 1 hour at 800 °C.

This calcination procedure (1 hour at 800-900 °C) should be performed on the porcelain crucible used for this operation.

4. Finally it is weighed at room temperature (keep it into a dessicator).

Determination of barium as BaSO₄. The precipitate is not carried out by slow addition of precipitant to the analyte containing solution. Such a procedure tends to produce conditions of super-saturation and a consequent rapid formation of small, relatively impure solid particles results. This effect cannot be entirely eliminated and is a source of error. The homogeneous precipitation techniques are able to avoid this erroneous effect; the precipitating reagent is generated slowly and homogeneously in presence of hydrochloric acid in boiling aqueous solution.

Procedure:

1. Determine mass of examined salt, using analytical balance.

Note on the $W_1 =$ _____ g (your result).

2. Weigh empty clean calcinated during 30 min. porcelain crucible **(do not forget number of crucible!)**

Note on the $W_{\text{crucible}} =$ _____ g (your result);

Number of crucible (or sign) _____.

3. Weighted mass of salt is transported to the 350-300 mL beaker, after then solid salt is dissolved by hot DW. Total volume of water in beaker must be not more 100-150 mL. Add 3 mL of 2N HCl for acidification of solution.

4. Prepare solution of precipitate agent in the other 100-150 mL beaker, added 4-5 mL of 2N H₂SO₄ and 25-50 mL of hot DW and stir up mixture.

5. Solutions of BaCl₂ and H₂SO₄ warm up practically to boiling state (to 80-90 °C). Very slowly (during at least 15 min.), by drops add solution of H₂SO₄ to solution of Barium salt, stirring well permanently by glass stick.

6. Let beaker with mixture settle for 24 hours at room temperature and protected from dust by paper list, **note on the surname of student.**

7. Test the fullness of precipitation adding 2-3 drops of 1N H₂SO₄ into beaker with precipitate. If no turbidity, it's OK.

8. Decant transpired upper layer (mother liquid) from beaker on paper filter, contained in funnel. Wash precipitate in beaker of 2-3 portions of hot DW (30-50 mL each portion).

9. Transport the precipitate from beaker to filter completely. Wash it 2-3 times of DW on filter.

10. Test the filtrate of absence of Cl^- with AgNO_3 . If your filtrate remains clear, dispose of the filtrate in the appropriate waste container.

8. Dry filter with precipitate at 110°C during 0,5-1 hour.

9. Ash filter with precipitate by heating in gas burner.

10. Then calcinate the precipitate with the aid of porcelain crucible in muffle at 800°C during 30-35 min.

11. Cool crucible to room temperature in a dessicator.

12. Weigh crucible with calcinated precipitate.

Note on the $W_2 =$ _____ g (your result).

13. Calculate content of barium in examined example of Barium chloride:

$$Z = \% \text{ Ba} = \frac{0,5884(W_2 - W_{\text{crucible}}) \times 100\%}{W_1} = \text{-----} = \text{-----} \%.$$

Check the result of the analysis in the teacher.

Calculate absolute and relative errors of analysis:

Absolute error:

$$\Delta = |\% \text{ Ba experimental (your result)} - \% \text{ Ba true (given by teacher)}| =$$

$$\text{-----} - \text{-----} = \text{-----}.$$

Relative error:

$$\sigma = \frac{\Delta}{\% \text{ Ba true}} \times 100\% = \text{-----}.$$

Remember, that relative error of gravimetry is not more 2%.

2.2. Laboratory work – Volumetric neutralization method. Determination of alkali concentration in solution

REAGENTS

1. Titrated solution of borax $\text{Na}_2\text{B}_4\text{O}_7$ (0,1000N);
2. Concentrated HCl solution (~38-40 w/w %);
3. Methyl orange indicator (solution);
4. DW.


GLASSWARE

1. 2-3 Erlenmeyer flasks, volume 250-300 mL;
2. 50 mL Burette fastened on a ring stand;
3. 20 mL Pipette;
4. 10 mL graduate cylinder;
5. Burette clamp;
6. Glass or plastic funnel (diameter 20-30 mm);
7. Wash bottle.

BACKGROUND

In the narrow sense in chemistry, neutralization means that equal amount of acids and bases in a solution react to produce salt and water. In the neutralization reaction between a strong acid and base in a solution, salt, which is produced, ionized almost perfectly so that hydrogen ions and hydroxide ions form water. Therefore, when acids and bases are equal, the hydrogen ion concentration ionizes to de-mineralized water (pH=7), which is neutral.

In chemical reactions, the various reacting substances always react in definite amounts. Taking the neutralization:



So, according to Law of equivalents 1 mole of HCl (36,5 g) will react exactly with 1 mole of NaOH (40,0 g).

Suppose we have two standard solutions: one of these (HCl) has a concentration of 1 M while the other (NaOH) also has a concentration of 1 M. Then, it follows that 1000 cm³ of that NaOH solution will react exactly with 1000 mL of the HCl solution, since these volumes contain **equimolar amounts** (the same number of moles) of the reactants. In fact, since definite amounts of acid and base are involved, we know that for the neutralization:

$$\begin{array}{l} \text{Acid} + \text{base} \rightarrow \text{salt} + \text{water}; \\ \text{Amount of acid} = \text{amount of base}; \\ \frac{\text{Volume of acid (mL)} \times \text{Normality of acid (g - eq/L)}}{1000} = \\ \frac{\text{Volume of base (mL)} \times \text{Normality of base (g - eq/L)}}{1000}. \end{array}$$

The above relationship is the basis for acid-base titrations. If we know one concentration accurately, as well as the two volumes, which neutralize each other exactly, we can easily calculate the unknown concentration.

Working solutions of neutralization method

Solutions of acids and alkalis are working solutions of method of neutralization. Solutions of H_2SO_4 , HNO_3 and NaOH , KOH are used more frequent. Dissolution of exact mass of the analytically pure substance (AR) in the definite volume of solution is the most exact method of preparation of titration working solutions. But it is practically impossible in the case of the indicated acids and alkalis, because they do not respond to requests to the initial matters and solutions and their exact initial concentration is unknown. Therefore working solutions are prepared with approximate concentration in the method of neutralization, and then their exact concentration is determined by initial solutions with the prepared concentration.

More frequent for determination of exact concentration of acids is used a **borax** $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$ or waterless salt $\text{Na}_2\text{B}_4\text{O}_7$ as a primary standard and oxalic acid $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$ is applied for determination of exact concentration of alkalis.

Preparation of HCl working solution. Order of preparation of solution:

- Take 4,5-5,0 mL concentrated Hydrochloric acid by the measured cylinder or conical measuring test tube;
- To carry it to glass jar on 0,5 L;
- To rinse a cylinder by DW and outpour this water in glass jar;
- To dilute acid in glass approximately to 0,5 L by DW;
- To close glass by a cork, to mix a reagent carefully;
- To sign the last name of student and name of reagent.

DETERMINATION OF NORMALITY OF HCl WORKING SOLUTION (Secondary standard)

Procedure:

1. Before the beginning of work to wash carefully all glassware.
2. To wash burette by HCl solution.
3. To wash pipette by borax solution.
4. Using funnel to fill a burette by HCl solution and obtain absence of air bubbles in the tap of burette.
5. To take off funnel and show out the level of liquid in a burette to the zero mark.
6. To take aliquot of borax solution by a pipette in washed by distilled water Erlenmeyer flask.
7. To add 1-2 drops of methyl orange indicator to borax solution in a flask; solution will becomes yellow.
8. To put a flask with borax solution under a burette and a sheet of white paper under flask.

9. To doze HCl solution slowly with constantly mixing a liquid in a retort by smooth circular motions.

10. It is necessary to catch a moment when yellow colour of solution in a flask will change on a pink from one drop of HCl.

11. To mark volume of HCl solution, used for titration.

12. To repeat titration of a new portion of borax solution.

TO MEMORIZE: any titration analyses must be conducted minimum three times (it is called parallel determinations). Thus the data about the volume of titrant which was used for titration are used for computations if they differ no more than on 0,1 mL. If three results have divergence more than 0,1 mL it is needed to conduct the fourth titration and etc. (until the absence of divergence).

Results fix in the table 1.

Table 1. Determination of accurate concentration of Hydrochloric acid solution used primary standard – borax

# of titration	Normality of borax solution, g-eq/L	Volume of borax in titration, mL (<i>equal to volume of pipette</i>)	Volume of HCl used for titration, mL	Normality of HCl, g-eq/L
1.	0,1000	20,0	<i>Your result</i>	0,____
2.	0,1000	20,0	<i>Your result</i>	
3.	0,1000	20,0	<i>Your result</i>	

$$N(\text{HCl}) = \frac{N(\text{borax}) \cdot V(\text{borax})}{V(\text{HCl for titration, average})} = \text{-----}$$

$$= 0, \text{---} \text{ g-eq/L.}$$

For calculation of normality of HCl solution applies average volume from two parallel titrations which differ no more than on 0,1 mL. The value of normality is calculated within **a fourth sign** after a comma. A result must be added to the table 1 (to calculate for the average value of the acid volume that was used for titration).

DETERMINATION OF ALKALI SOLUTION NORMALITY

REAGENTS

1. HCl solution with definite normal concentration (see table 1);
2. Methyl orange indicator.

GLASSWARE: alike in a previous section.

Procedure: it is similar to the previous section, but instead of borax solution must be taken alkali solution. Results write in the table 2.

Table 2. Determination of equivalent concentration of NaOH

# of titration	Normality of HCl solution, g-eq/L	Volume of HCl in titration, mL	Volume of NaOH for determination, mL (equal to volume of pipette)	Normality of NaOH, g-eq/L
1.	From previous table, last column	<i>Your result</i>	20,0	0, _ _ _ _
2.		<i>Your result</i>	20,0	
3.		<i>Your result</i>	20,0	

$$N(\text{NaOH}) = \frac{N(\text{HCl}) \cdot V(\text{HCl for titration, average})}{V(\text{NaOH})} = \text{-----} =$$

0, _ _ _ _ g-eq/L.

Check the result of the analysis in the teacher.

Calculate absolute and relative errors of analysis:

Absolute error:

$$\Delta = |N(\text{NaOH})_{\text{experimental}} (\text{your result}) - N(\text{NaOH})_{\text{true}} (\text{given by teacher})|$$

$$= \text{-----} - \text{-----} = \text{-----}.$$

Relative error:

$$\sigma = \frac{\Delta}{N(\text{NaOH true})} \times 100\% = \text{-----}.$$

Remember, that relative error of volumetry is not more 1%.

2.3. Lab work – Oxidation-reduction (RedOx) titration (RedOxmetry). Determination of Iron(II) content in Mohr's salt



REAGENTS

1. Primary standard of oxalate acid $\text{H}_2\text{C}_2\text{O}_4$ solution (0,0500 N);
2. Working solution Potassium Permanganate KMnO_4 , approximately 0,05N;
3. 2N Sulfate acid H_2SO_4 solution;
4. DW.

GLASSWARE

1. 2-3 Erlenmeyer (titration conical) flasks, volume 250-300 mL;
2. 50 mL Burette fastened on a ring stand;
3. 20 mL Pipette;
4. Laboratory tilt measure (10 mL);
5. Burette clamp;
6. Lab electric oven;
7. Glass or plastic funnel (diameter 20-30 mm);
8. Wash bottle.

**DETERMINATION OF CONCENTRATION OF KMnO_4
WORKING SOLUTION**

BACHGROUND**Reaction based on method:**

If precise concentration of oxalate acid is known and reacted volumes of permanganate and oxalate acid solutions, the precise concentration of KMnO_4 may be calculated according Law of Equivalents:

$$N(\text{KMnO}_4) = \frac{N(\text{H}_2\text{C}_2\text{O}_4) \cdot V(\text{H}_2\text{C}_2\text{O}_4)}{V(\text{KMnO}_4)}.$$

Procedure

1. Clean pipette, burette and Erlenmeyer flasks, as described above.
2. Add 20 mL of oxalate acid solution from measuring pipette to 250-300 mL Erlenmeyer flask. Acidify this solution adding 20 mL of 2N H_2SO_4 using tilt measure. Add near 40 mL of DW. Heat prepared solution to 70-80 °C. **BE CAREFUL, NOT BOIL SOLUTION** (because oxalic acid decomposes at 100 °C)!
3. Using a burette, slowly add the KMnO_4 solution to hot solution in flask. KMnO_4 acts as its own indicator. That is, at the start of the titration the deep violet colour of MnO_4^- will be lost because it is changing to Mn^{2+} , but as soon as there is no more oxalate acid in the solution for the MnO_4^- to react with, the MnO_4^- will remain in the reaction solution. The slightly pink colour of the dilute solution MnO_4^- indicates the end of the reaction. This colour must safe during at least 1 min. This is known as the "end point" or equivalence point of the titration.

4. The procedure of titration repeats two times or more. The results of so-called parallel titrations must distinguish one from other not more than 0,1 mL .

Results of experiment present in the form of table 3.

Table 3. Determination of accurate concentration of Permanganate solution used primary standard – oxalate acid

# of titration	Normality of oxalate acid $\text{H}_2\text{C}_2\text{O}_4$ solution, g-eq/L	Volume of oxalate acid $\text{H}_2\text{C}_2\text{O}_4$ solution in titration, mL (<i>volume of pipette</i>)	Volume of KMnO_4 used for titration, mL	Normality of KMnO_4 , g-eq/L
1.	0,0500	20,0	<i>Your result</i>	0,____ _ _ _
2.	0,0500	20,0	<i>Your result</i>	
3.	0,0500	20,0	<i>Your result</i>	



Accurate concentration of permanganate solution is equal to:

$$N(\text{KMnO}_4) = \frac{N(\text{H}_2\text{C}_2\text{O}_4) \cdot V(\text{H}_2\text{C}_2\text{O}_4)}{V(\text{KMnO}_4)} = =$$

$$= \frac{0,0500 \text{ g - eq/L} \cdot 20 \text{ mL}}{\text{your result} \text{ mL}} = 0, _ _ _ _ \text{ g - eq/L.}$$

DETERMINATION OF IRON(II) CONCENTRATION IN THE MOHR'S SALT SOLUTION

Mohr's salt is a double salt - Iron(II)-Ammonium sulphate hexahydrate ($\text{Fe}(\text{NH}_4)_2(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$) included ions of Fe(II). The last one may reacts with potassium permanganate in acidic medium:



This reaction isn't heating, because ion Fe (II) may oxidizes by air oxygen transformed to Fe (III) with rising of temperature.

Procedure

1. Add 20 mL of Mohr's salt solution (**according to your variant of task**) with unknown concentration used measuring pipette to 250-300 mL Erlenmeyer flask. Acidify this solution adding 20 mL of 2N H_2SO_4 using

graduated cylinder. Add near 40 mL of DW. **BE CAREFUL, DON'T HEAT SOLUTION!**

2. Using a burette, slowly add the KMnO_4 solution to hot solution in flask. The slightly pink colour of the dilute solution MnO_4^- (stable 1 min. at least) indicates the end of the reaction. This colour must safe during at least 1 min. The procedure of titration repeats two times or more as described above.

Results of experiment present in the form of table 4:

Table 4. Determination of Iron(II) concentration in Mohr's salt solution

# of titration	Normality of KMnO_4 solution, g-eq/L	Volume of KMnO_4 in titration, mL	Volume of Mohr's salt solution for determination, mL (<i>volume of pipette</i>)	Equivalent concentration (normality) of Fe^{2+} in examined solution, g-eq/L
1.	From previous table, last column 0, _ _ _ _	<i>Your result</i>	20,0	0, _ _ _ _
2.		<i>Your result</i>	20,0	
3.		<i>Your result</i>	20,0	

Calculate the concentration of Fe^{+2} present in the Mohr's salt solution:

$$N(\text{Fe}^{2+}) = \frac{N(\text{KMnO}_4) \cdot V(\text{KMnO}_4 \text{ avarage})}{V(\text{Mohr's salt solution})} = \frac{\quad}{20\text{mL}} =$$

$$= 0, _ _ _ _ \text{ g-eq/L.}$$

Check the result of the analysis in the teacher.

Calculate absolute and relative errors of analysis:

Absolute error:

$$\Delta = N(\text{Fe}^{2+} \text{ experimental (your result)}) - N(\text{Fe}^{2+} \text{ true (given by teacher)}) =$$

$$\underline{\quad} - \underline{\quad} = \underline{\quad}.$$

Relative error:

$$\sigma = \frac{\Delta}{N(\text{Fe}^{2+} \text{ true})} \times 100\% = \underline{\quad}.$$

Remember, that relative error of volumetry is not more 1%.

2.4. Laboratory work – Iodometry. Determination of Copper(II) Content in Copper Vitriol

REAGENTS

1. 0,0500N Standard Thiosulfate $\text{Na}_2\text{S}_2\text{O}_3$;
2. 2N acetate acid CH_3COOH solution;
3. 10% Potassium Iodide KI solution;
4. Starch, fresh-prepared solution;
5. DW.

GLASSWARE


1. Watch glass, diameter 70-80 mm;
2. 100 mL Measuring flask;
2. 2-3 Erlenmeyer flasks, volume 250-300 mL;
3. 50 mL Burette fastened on a ring stand;
4. 20 mL Pipette;
5. Laboratory tilt measure (10 mL);
6. 5 mL cylinder;
7. Burette clamp;
8. Spatula;
9. Glass or plastic funnel (diameter 20-30 mm);
10. Wash bottle.

EQUIPMENT


Analytical balance.

BACKGROUND



Iodometric determination of copper is based on the oxidation of iodides to iodine by copper (II) ions, which get reduced to Cu^+ . Reaction taking place in the solution is:



and produced equivalent amount of iodine can be titrated with Thiosulfate solution:



Finally, united process may be presented as:



For the best results reaction should take place in the slightly acidic solution (pH near 4-5), adding a few drops of acetate acid. Solution should be free of other substances that can oxidize iodides to iodine (for example, Fe^{3+} or nitrites).

Procedure

1. Weight out accurately near 1,3 g of the analyzed salt.

Note on the $W_1 = 1, \text{---} \text{---} \text{---} \text{ g}$.

2. Transport the probe in 100 mL measuring flask, dissolve in DW. Add 10-15 drops of 2N Acetate acid CH_3COOH . Make up volume of solution to 100 mL. Shake well.

3. Pipette 20,0 mL of this solution into a 250 mL titration conical flask, add 10 mL of 10% KI solution (using tilt measure);

4. Place the conic flask into dark place for 5-10 min;

5. Titrate swirling the flask the liberated iodine with standard 0,0500N sodium thiosulphate until a pale yellow colour of mixture;

6. Add 1 mL of fresh prepared starch solution – indicator;

7. Titrate swirling the flask, until blue colour disappears.

6. Repeat the titration. Results write in table 5.

Table 5. Determination of Copper(II) content in Copper Vitriol

# of titration	Normality of $\text{Na}_2\text{S}_2\text{O}_3$ solution, g-eq/L	Volume of $\text{Na}_2\text{S}_2\text{O}_3$ in titration, mL	Volume of copper(II) solution for determination, mL (<i>volume of pipette</i>)	Copper(II) content in Copper Vitriol, wt. %
1.	0,0500	<i>Your result</i>	20,0	_____, ____
2.	0,0500	<i>Your result</i>	20,0	
3.	0,0500	<i>Your result</i>	20,0	

$$\text{Wt. \% Cu} = \frac{N(\text{Na}_2\text{S}_2\text{O}_3) \cdot V(\text{Na}_2\text{S}_2\text{O}_3 \text{ for titration, average}) \cdot E(\text{Cu}) \cdot V(\text{flack}) \cdot 100\%}{V(\text{pipette}) \cdot 1000 \cdot W_1} =$$

$$\frac{0,0500 \cdot V(\text{Na}_2\text{S}_2\text{O}_3 \text{ for titration, average}) \cdot 63,543 \cdot 100 \cdot 100}{20,0 \cdot 1000 \cdot W_1}$$

$$= \frac{1,58858 \cdot V(\text{Na}_2\text{S}_2\text{O}_3 \text{ for titration})}{W_1} = \frac{1,58858 \cdot \text{---}}{\text{---}} \text{---}, \text{---} \%$$

Check the result of the analysis in the teacher.

Calculate absolute and relative errors of analysis:

Absolute error:

$$\Delta = |\% \text{Cu}_{\text{experimental}} (\text{your result}) - \% \text{Cu}_{\text{true}} (\text{given by teacher})| =$$

$$= \text{---} - \text{---} = \text{---}.$$

Relative error:

$$\sigma = \frac{\Delta}{\% \text{Cu true}} \times 100\% = \text{---} \%$$

Remember, that relative error of volumetry is not more 1%.

2.5. Laboratory work – Complexonometry. Determination of Calcium concentration in solution

REAGENTS

1. Primary standard -0,0500N Zinc Sulfate solution $ZnSO_4$;
2. Working solution Trilon B, approximately 0,05N;
3. Ammonia buffer solution (pH=10);
4. Eriochrome black T indicator, solid, 1% (w/w) in Sodium Chloride;
5. DW.


GLASSWARE

1. 2-3 Erlenmeyer flasks, volume 250-300 mL;
2. 50 mL Burette fastened on a ring stand;
3. 20 mL Pipette;
4. 10 mL cylinder;
5. Burette clamp;
6. Glass or plastic funnel (diameter 20-30 mm);
7. Wash bottle.

BACKGROUND

The most important one in titrimetric analysis has a complexing of metals ions with halide- or pseudo-halide ions (SCN^- , CN^-), and with group of polyamine carboxylic acids (chelators).


The most commonly used reagent for this method Trilon B may forms a very stable complex compound with a great number of cations (alkali-earth metals, copper, zinc, iron, manganese etc.) according to the reaction:



Different cations form stable complexes with Trilon B depending on their nature. For example, triple-charged cations form stable complexes when pH very low (1-3), while cations of alkaline earth metals – at pH 10.

Determination of titration endpoint is realized using special metallochromic indicators. They are a weak complexing agents made own

colour and formed colour complexing compounds with metal cations. Last ones have a weak stability. One of such indicators is an Eriochrome black T. Its own colour in solution - sky-blue, when colour of his complexes with metals – wine-red. In general, schemes of titration processes are:



DETERMINATION OF NORMALITY OF TRILON B WORKING SOLUTION (Secondary standard)

Procedure:

1. Before the beginning of work to wash carefully all glassware.
2. To wash burette by Trilon B solution.
3. To wash pipette by Zinc Sulfate ZnSO_4 solution.
4. Using funnel to fill a burette by Trilon B solution and to remove of air bubbles from the tap of burette.
5. To take off funnel and show out the level of liquid in a burette to the zero mark.
6. To take aliquot of Zinc Sulfate ZnSO_4 solution by a pipette in washed by DW Erlenmeyer flask.
7. To add 5 mL Ammonia buffer (use 10 mL cylinder), and a few grains of Eriochrome black T indicator to Zinc Sulfate ZnSO_4 solution in a flask; solution will becomes red-wine.
9. To titrate adding Trilon B solution slowly swirling the flask.
10. To fix volume of Trilon B used for titration, in the moment when colour of mixture in the flask changes from wine-red to blue after adding of one drop of titrant.
11. To repeat titration for a new portion of Zinc Sulfate ZnSO_4 solution.

Results of experiment present in the form of table 6:

Table 6. Determination of accurate concentration of Trilon B solution used primary standard – Zinc Sulfate ZnSO₄

# of titration	Normality of Zinc Sulfate ZnSO ₄ , g-eq/L	Volume of ZnSO ₄ in titration, mL (equal to volume of pipette)	Volume of Trilon B used for titration, mL	Normality of Trilon B, g-eq/L
1.	0,_____	20,0	<i>Your result</i>	0,_____
2.	0,_____	20,0	<i>Your result</i>	
3.	0,_____	20,0	<i>Your result</i>	

$$N(\text{Trilon B}) = \frac{N(\text{ZnSO}_4) \cdot V(\text{ZnSO}_4)}{V(\text{Trilon B for titration, average})} =$$

$$\frac{\text{_____} \cdot 20,0}{\text{_____}} =$$

$$= 0, \text{---} \text{---} \text{---} \text{---} \text{ g-eq/L.}$$

DETERMINATION OF NORMALITY OF CALCIUM Ca²⁺ SOLUTION

REAGENTS

1. Trilon B solution with definite normal concentration (see table 6);
2. Ammonia buffer solution (pH=10);
3. Eriochrome black T indicator, solid, 1% (w/w) in Sodium Chloride;
4. DW.

GLASSWARE: alike in a previous section.

Procedure

It is similar to the previous section, but instead of Zinc Sulfate ZnSO₄ solution must be taken Calcium Ca²⁺ solution.

Results write in the table 7.

Table 7. Determination of Calcium Ca^{2+} concentration in solution

# of titration	Normality of Trilon B solution, g-eq/L	Volume of Trilon B in titration, mL	Volume of Calcium solution for determination, mL (equal to volume of pipette)	Normality of Ca^{2+} of solution, g-eq/L
1.	From previous	<i>Your result</i>	20,0	0, _ _ _ _
2.	table, last column	<i>Your result</i>	20,0	
3.	0, _ _ _ _	<i>Your result</i>	20,0	

$$N(\text{Ca}^{2+}) = \frac{N(\text{Trilon B}) \cdot V(\text{Trilon B for titration, average})}{V(\text{Ca solution})} =$$

$$\frac{\cdot}{20,0} =$$

$$= 0, _ _ _ _ \text{ g-eq/L.}$$

Check the result of the analysis in the teacher.

Calculate absolute and relative errors of analysis:

Absolute error:

$$\Delta = |N(\text{Ca}^{2+})_{\text{experimental}} (\text{your result}) - N(\text{Ca}^{2+})_{\text{true}} (\text{given by teacher})| =$$

$$= _ _ _ - _ _ _ = _ _ _.$$

Relative error:

$$\sigma = \frac{\Delta}{N(\text{Ca}^{2+} \text{ true})} \times 100\% = _ _ _ _.$$

Remember, that relative error of volumetry is not more 1%.

EXAMPLES OF CONTROL TESTS FOR QUANTITATIVE ANALYSIS

1. Calculate ionic strength I of solution contained 17,43 g K_2SO_4 ($M=174,26$ g/mol) per 1 L.

A	0, 10 mol/L	B	0,20 mol/L	C	0,30 mol/L	D	0,50 mol/L
----------	-------------	----------	------------	----------	------------	----------	------------

2. Calculate gravimetric factor f for the quantitative determination of Magnesium content in the form MgO ($M=40,31$ g/mol), if the weighted form of the sediment is $Mg_2P_2O_7$ ($M=222,57$ g/mol).

3. Note on the absorption indicator for RedOx volumetric method:

A	Eriochrome black T	C	Methyl orange
B	Starch	D	Phenolphthalein

4. Calculate ppt (‰) concentration of solution contains 0,01 g of solute per 50 g of solution.

A	0, 200	B	1,000	C	0,500	D	0,001
----------	--------	----------	-------	----------	-------	----------	-------

5. Calculate pH of 10 N hydrochloride acid HCl (to consider as a strong acid, dissociated completely):


A	1	B	10	C	0	D	-1
----------	---	----------	----	----------	---	----------	----

6. Put a missing word: *A standard solution is a solution containing a precisely known _____ of an element or a substance i.e., a known _____ of solute is dissolved to make a specific volume.*


7. Calculate mass of $FeSO_4 \cdot 7H_2O$ ($MW=278,01$ g/mol) for the Iron gravimetric determination on the form of Fe_2S_3 ($MW=207,89$ g/mol) according to the Tananaev's recommendation.

8. Sample of lime ($m_{\text{sample}} = 0,8960$ g) was dissolved in the nitrate acid and total volume of obtained solution was 200 mL ($V_{\text{measuring flask}}$). Aliquot of this solution (20 mL, V_{pipette}) was titrated triply by 0,0505N Trilon B and obtained the next results: 14,6; 13,6; 13,5 mL. Calculate Calcium content (w/w %) in this sample.

9. Figure below demonstrates the curve of titration of _____ (weak or strong) _____ (acid or base) by _____ (weak or strong) _____ (acid or base) (put the missing words).



10. Refer to the titration curve below for questions:



- The titration curve shown above describes the:
 - A. Titration of a strong base.
 - B. Titration of a weak acid.
 - C. Titration of a strong acid.
 - D. Titration of a weak base.
- The equivalence point occurs at:
 - A. 7,50 mL titrant added.
 - B. 30,0 mL titrant added.
 - C. 15,0 mL titrant added.
 - D. 22,5 mL titrant added.
- The titrant is a
 - A. Strong acid.
 - B. Strong base.
 - C. Weak acid.
 - D. Weak base.

11. Note on the figure of Fixanale :



A



B



C




D



E

12. What is aim of the next previous operation of titration (see picture)?



13. Calculate the pH of a 0,1 M Na_2CO_3 ($\text{pK}_1 = 6,52$; $\text{pK}_2 = 10,22$) and 0,1 M NaOH.

14. Determine the types of sediments in gravimetric analysis:

A	Crystalline	1	CoS
B	Amorphous	2	SrSO ₄
		3	Fe(OH) ₂
		4	MgNH ₄ PO ₄

15. Determine the active concentrations ($a = \gamma \cdot c$) of 0,01 M Cl^- solution and 0,0001 M Al^{3+} solution in soil solution where ionic power of solution is equals to 0,0005 (See APPENDIX).

APPENDIX Ionic activity coefficients at different ionic strength of solution


Ionic strength I, mol/L	Activity coefficients γ for ions with charge Z_i			
	1+ or 1-	2+ or 2-	3+ or 3-	4+ or 4-
0,0001	0,99	0,95	0,90	0,83
0,0002	0,98	0,94	0,87	0,77
0,0005	0,97	0,90	0,80	0,67
0,001	0,96	0,86	0,73	0,56
0,002	0,95	0,81	0,64	0,45
0,005	0,92	0,72	0,51	0,30
0,01	0,89	0,63	0,39	0,18
0,02	0,87	0,57	0,28	0,12
0,05	0,84	0,50	0,21	0,06
0,1	0,81	0,44	0,16	0,04

16. What is the solubility (M) of SrSO_4 in a solution that already contains 0.0600 M SO_4^{2-} ? The K_{sp} for SrSO_4 is $3,2 \cdot 10^{-7}$ (Effect of common ion).

17. Calculate equivalent weight of a $\text{K}_2\text{Cr}_2\text{O}_7$ ($M=294,10$ g/mol) for RedOx titration in acidic medium.

18. Calculate equivalent weight of a H_3PO_4 ($M=97,99$ g/mol) for neutralization reaction: $\text{H}_3\text{PO}_4 + \text{NaOH} = \text{NaH}_2\text{PO}_4 + \text{H}_2\text{O}$.

19. Calculate volume (in mL) of 50% Sulfate acid solution (density $d=1,40 \text{ g/cm}^3$) for preparation of 5 L 0,01 N solution (Molar weight (H_2SO_4) = 98 g/mol).
20. Calculate molarity of HCl solution of the Titre 0,01 g/mL.
21. Calculate pH of a 0,1N NH_4Cl solution ($\text{pK}(\text{NH}_3 \cdot \text{H}_2\text{O})=4,77$)
22. Calculate pH of a 10N hydrochloric acid HCl (to consider as a strong acid, dissociated completely)
23. Calculate the pH of a 0,1 M H_3PO_4 ($\text{pK}_1=1,96$; $\text{pK}_2=6,70$; $\text{pK}_3=12,44$).
24. Calculate the pH of a 0,01 M NaClO ($\text{pK}_{\text{acid}} = 7,50$).
25. Calculate the pH of a 1 M Na_2CO_3 ($\text{pK}_1 = 6,52$; $\text{pK}_2 = 10,22$).
26. Calculate ppm concentration of solution contains 0,05 g of solute per 1000 g of solution.
27. Calculate solubility (in mol/L and g/L) of $\text{Ni}(\text{OH})_2$ ($\text{SP}=1,6 \cdot 10^{-14}$, $\text{MW}=92,71$ g/mol) in solution of 0,1N KOH (*Be attentive: effect of the common ion into solubility*).
28. Calculate Copper ($\text{AW}=63,548$ g/mol) percent content in 1,0000 g malachite sample, if for iodometric titration 20 mL solution taken of 200 mL measuring flask was used 15,0 mL 0,0500N $\text{Na}_2\text{S}_2\text{O}_3$.
29. Note on the true result of titration (see a figure below)



30. Calculate the RedOx system potential E, included 0,1 M FeCl_3 and 0,01 M FeCl_2 ($E^\circ(\text{Fe}^{3+}/\text{Fe}^{2+}) = +0,771 \text{ V}$).
31. Calculate the RedOx system potential E, included 0,1 M KMnO_4 and 0,01 M MnSO_4 at $\text{pH}=1$ ($E^\circ(\text{MnO}_4^-/\text{Mn}^{2+}) = +1,51 \text{ V}$).
32. Calculate E of the RedOx system, included zinc electrode, immersed into 1 M ZnSO_4 if $E^\circ(\text{Zn}^{2+}/\text{Zn}^0) = -0,764 \text{ V}$.

33. Equivalent mass of Ce^{3+} cation (AW=140,12 g/mol) in reaction with Trilon B is, g-eq/g:

A	70,06	B	46,83	C	140,12	D	187,32
----------	-------	----------	-------	----------	--------	----------	--------

34. Determine the order of steps of iodometric analysis.

A	To add excess of KI to analysing solution
B	To add indicator starch
C	To titrate by thiosulphate solution to slightly yellow colour of solution
D	To stand flask with analysing solution and a certain reactant in dark place (near 15 min).
E	To titrate by thiosulphate solution to decolourizing of solution

35. Calculate theoretical percent content (w/w) of Magnesium (AW=24,31 g/mol) in epsomite $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (MW= 246,48 g/mol).

MY RATING TABLE

THEORETICAL QUIZZES	Points	EXPERIMENTAL TASKS	Points
Module test 1. Qualitative analysis of cations		1. Analysis of cations mixture.	
Module test 2. Qualitative analysis of substances		2. Analysis of a soluble salt	
		3. Analysis of insoluble substance	
Module test 3. Heterogeneous equilibrium. Gravimetry		4. Gravimetric determination of Barium content in Hydrated Barium Chloride	
Module test 4. Homogeneous equilibrium. Volumetry.		5. Volumetric determination of alkali solution normality	
		6. Permanganatometric determination of Iron(II) concentration in the solution of Mohr's salt	
		7. Iodometric determination of Copper(II) content in Copper Vitriol	
		8. Complexometric determination of normality of Calcium Ca^{2+} solution	